

Anti-GPR158 Antibody

Rabbit polyclonal antibody to GPR158

Catalog # AP60463

Product Information

Application	WB
Primary Accession	Q5T848
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	135489

Additional Information

Gene ID	57512
Other Names	KIAA1136; Probable G-protein coupled receptor 158
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human GPR158. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	GPR158 {ECO:0000303 Ref.1, ECO:0000312 HGNC:HGNC:23689}
Function	<p>Metabotropic receptor for glycine that controls synapse formation and function in the brain (PubMed:36996198). Acts as an atypical G-protein coupled receptor that recruits and regulates the RGS7-GNB5 complex instead of activating G proteins (PubMed:31189666, PubMed:36996198). In absence of glycine ligand, promotes the GTPase activator activity of RGS7, increasing the GTPase activity of G protein alpha subunits, thereby driving them into their inactive GDP-bound form (PubMed:36996198). Glycine-binding changes the conformation of the intracellular surface, inhibiting the GTPase activator activity of the RGS7-GNB5 complex, promoting G protein alpha subunits into their active GTP-bound form and regulating cAMP levels (PubMed:36996198). Also able to bind taurine, a compound closely related to glycine, but with a two- fold lower affinity (PubMed:36996198). Glycine receptor-dependent regulation of cAMP controls key ion channels, kinases and neurotrophic factors involved in neuronal excitability and synaptic transmission (PubMed:36996198). Plays a pivotal role in regulating mood and cognition via</p>

its ability to regulate neuronal excitability in L2/L3 pyramidal neurons of the prefrontal cortex (By similarity). Also involved in spatial learning by regulating hippocampal CA1 neuronal excitability (By similarity). Acts as a synaptic organizer in the hippocampus, required for proper mossy fiber-CA3 neurocircuitry establishment, structure and function: induces presynaptic differentiation in contacting axons via its interaction with GPC4 (By similarity). In addition to glycine, may also act as a receptor for osteocalcin (BGLAP) hormone: osteocalcin-binding initiates a signaling response that prevents neuronal apoptosis in the hippocampus and regulates the synthesis of neurotransmitters (By similarity).


Cellular Location

Cell membrane; Multi-pass membrane protein. Postsynaptic cell membrane {ECO:0000250|UniProtKB:Q8C419}; Multi-pass membrane protein. Presynaptic cell membrane {ECO:0000250|UniProtKB:Q8C419}; Multi-pass membrane protein Nucleus Note=Mainly localizes to the postsynaptic membrane, with a small portion to the presynaptic membrane (By similarity). Trafficks between the nucleus and the cell membrane; it is unclear how a multi-pass membrane protein can traffick between the nucleus and the cell membrane (PubMed:23451275). {ECO:0000250|UniProtKB:Q8C419, ECO:0000269|PubMed:23451275}

Background

KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human GPR158. The exact sequence is proprietary.

Images



Western blot analysis of GPR158 expression in BV2 (A), PMVEC (B), A549 (C), HepG2 (D) whole cell lysates.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.