

Anti-KDELR2 Antibody

Rabbit polyclonal antibody to KDELR2
Catalog # AP60475

Product Information

Application	WB, IF/IC, IHC
Primary Accession	P33947
Other Accession	Q9CQM2
Reactivity	Human, Mouse, Rat, Zebrafish, Pig, Chicken, Bovine, SARS
Host	Rabbit
Clonality	Polyclonal
Calculated MW	24422

Additional Information

Gene ID	11014
Other Names	ERD2.2; ER lumen protein retaining receptor 2; ERD2-like protein 1; ELP-1; KDEL endoplasmic reticulum protein retention receptor 2; KDEL receptor 2
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human KDELR2. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500) IF/IC~~N/A IHC~~WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C. Stable for 12 months from date of receipt

Protein Information


Name	KDELR2
Synonyms	ERD2.2 {ECO:0000303 PubMed:1325562}
Function	Membrane receptor that binds the K-D-E-L sequence motif in the C-terminal part of endoplasmic reticulum resident proteins and maintains their localization in that compartment by participating to their vesicle-mediated recycling back from the Golgi (PubMed: 1325562 , PubMed: 18086916 , PubMed: 33053334). Binding is pH dependent, and is optimal at pH 5-5.4 (By similarity).
Cellular Location	Endoplasmic reticulum membrane; Multi-pass membrane protein {ECO:0000250 UniProtKB:Q5ZKX9}. Golgi apparatus membrane; Multi-pass

membrane protein {ECO:0000250 | UniProtKB:Q5ZKX9}. Cytoplasmic vesicle, COPI-coated vesicle membrane; Multi-pass membrane protein {ECO:0000250 | UniProtKB:Q5ZKX9} Note=Localized in the Golgi in the absence of bound proteins with the sequence motif K-D-E-L. Trafficks back to the endoplasmic reticulum together with cargo proteins containing the sequence motif K-D-E-L


Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human KDELR2. The exact sequence is proprietary.


Images



Western blot analysis of KDELR2 expression in rat testis (A) whole cell lysates.



Immunohistochemical analysis of KDELR2 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of KDELR2 staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.