

Anti-TOPK (pT9) Antibody

Rabbit polyclonal antibody to TOPK (pT9)

Catalog # AP60497

Product Information

Application	WB, IHC
Primary Accession	Q96KB5
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	36085

Additional Information

Gene ID	55872
Other Names	TOPK; Lymphokine-activated killer T-cell-originated protein kinase; Cancer/testis antigen 84; CT84; MAPKK-like protein kinase; Nori-3; PDZ-binding kinase; Spermatogenesis-related protein kinase; SPK; T-LAK cell-originated protein kinase
Target/Specificity	Recognizes endogenous levels of TOPK (pT9) protein.
Dilution	WB~~WB (1/500 - 1/1000), IHC (1/100 - 1/200) IHC~~WB (1/500 - 1/1000), IHC (1/100 - 1/200)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt


Protein Information

Name	PBK
Synonyms	TOPK
Function	Phosphorylates MAP kinase p38. Seems to be active only in mitosis. May also play a role in the activation of lymphoid cells. When phosphorylated, forms a complex with TP53, leading to TP53 destabilization and attenuation of G2/M checkpoint during doxorubicin- induced DNA damage.
Tissue Location	Expressed in the testis and placenta. In the testis, restrictedly expressed in outer cell layer of seminiferous tubules.


Background

KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human TOPK (pT9). The exact sequence is proprietary.

Images



Western blot analysis of TOPK (pT9) expression in Myla2059 (A), SHSY5Y (B) whole cell lysates.



Immunohistochemical analysis of TOPK (pT9) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.