

# Anti-CDCA4 Antibody

Rabbit polyclonal antibody to CDCA4

Catalog # AP60550

## Product Information

---

<b>Application</b>	WB, IF/IC, IHC
<b>Primary Accession</b>	<a href="#">Q9BXL8</a>
<b>Reactivity</b>	Human, Rat
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	26114

## Additional Information

---

<b>Gene ID</b>	55038
<b>Other Names</b>	HEPP; Cell division cycle-associated protein 4; Hematopoietic progenitor protein
<b>Target/Specificity</b>	Recognizes endogenous levels of CDCA4 protein.
<b>Dilution</b>	WB~~WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500) IF/IC~~N/A IHC~~WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500)
<b>Format</b>	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
<b>Storage</b>	Store at -20 °C.Stable for 12 months from date of receipt

## Protein Information


---

<b>Name</b>	CDCA4
<b>Synonyms</b>	HEPP
<b>Function</b>	May participate in the regulation of cell proliferation through the E2F/RB pathway. May be involved in molecular regulation of hematopoietic stem cells and progenitor cell lineage commitment and differentiation (By similarity).
<b>Cellular Location</b>	Nucleus.
<b>Tissue Location</b>	Highest levels of expression in the pancreas, thymus, testis, spleen, liver, placenta and leukocytes. Relatively low levels in the lung, kidney, prostate, ovary, small intestine and colon Hardly detectable, if at all, in the brain, skeletal muscle and heart


## Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human CDCA4. The exact sequence is proprietary.


## Images



Western blot analysis of CDCA4 expression in HEK293T (A), PC3 (B), A549 (C) whole cell lysates.



Immunohistochemical analysis of CDCA4 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of CDCA4 staining in HEK293T cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.