

Anti-Histone H2A.Z (AcK5) Antibody

Rabbit polyclonal antibody to Histone H2A.Z (AcK5)

Catalog # AP60570

Product Information

Application	WB, IF/IC, IHC
Primary Accession	P0C0S5
Other Accession	P0C0S6
Reactivity	Human, Mouse, Rat, Zebrafish, Monkey, Pig, Bovine, Dog, SARS
Host	Rabbit
Clonality	Polyclonal
Calculated MW	13553

Additional Information

Gene ID	3015
Other Names	H2AZ; Histone H2A.Z; H2A/z
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human Histone H2A.Z. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500) IF/IC~~N/A IHC~~WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt


Protein Information

Name	H2AZ1 (HGNC:4741)
Function	Variant histone H2A which replaces conventional H2A in a subset of nucleosomes. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post- translational modifications of histones, also called histone code, and nucleosome remodeling. May be involved in the formation of constitutive heterochromatin. May be required for chromosome segregation during cell division.
Cellular Location	Nucleus. Chromosome.


Background

KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human Histone H2A.Z. The exact sequence is proprietary.


Images



Western blot analysis of Histone H2A.Z (AcK4) expression in HeLa (A), COS7 (B), MEF (C) whole cell lysates.



Immunohistochemical analysis of Histone H2A.Z (AcK5) staining in human colon formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of Histone H2A.Z (AcK5) staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.