10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-NSG1 Antibody

Catalog # AP60601

Product Information

Application	WB, IHC, IF/IC
Primary Accession	P42857
Reactivity	Human, Mouse, Rat, Mk
Host	Rabbit
Clonality	Polyclonal
Calculated MW	20913

Additional Information


Gene ID	27065
Other Names	D4S234; Neuron-specific protein family member 1; Brain neuron cytoplasmic protein 1
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human NSG1. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000), IHC (1/100 - 1/200) IHC~~WB (1/500 - 1/1000), IHC (1/100 - 1/200) IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	NSG1 (HGNC:18790)
Function	Plays a role in the recycling mechanism in neurons of multiple receptors, including AMPAR, APP and L1CAM and acts at the level of early endosomes to promote sorting of receptors toward a recycling pathway. Regulates sorting and recycling of GRIA2 through interaction with GRIP1 and then contributes to the regulation of synaptic transmission and plasticity by affecting the recycling and targeting of AMPA receptors to the synapse (By similarity). Is required for faithful sorting of L1CAM to axons by facilitating trafficking from somatodendritic early endosome or the recycling endosome (By similarity). In an other hand, induces apoptosis via the activation of CASP3 in response to DNA damage (PubMed:20599942, PubMed:20878061).
Cellular Location	Membrane {ECO:0000250|UniProtKB:P02683}; Single- pass type II membrane protein {ECO:0000250|UniProtKB:P02683}. Golgi apparatus, trans-Golgi network membrane {ECO:0000250|UniProtKB:P02683} Endosome membrane

{ECO:0000250|UniProtKB:P02683}. Cell projection, dendrite {ECO:0000250|UniProtKB:P02683}. Early endosome membrane {ECO:0000250|UniProtKB:P02683}. Late endosome membrane {ECO:0000250|UniProtKB:P02683}. Lysosome lumen {ECO:0000250|UniProtKB:P02683}. Recycling endosome membrane {ECO:0000250|UniProtKB:P02683}. Cytoplasmic vesicle membrane {ECO:0000250|UniProtKB:P02683}. Golgi apparatus, Golgi stack membrane {ECO:0000250|UniProtKB:P02683}. Endosome, multivesicular body membrane {ECO:0000250|UniProtKB:P02683}. Endoplasmic reticulum membrane. Note=Endocytosed from the cell surface, thus enters into early endosomes, trafficks to late endosomes and degradates in lysosomes (By similarity). Endoplasmic reticulum targeting is essential for apoptosis (PubMed:20599942). Found in both stationary and motile endosomes. A previous study supports a type I membrane protein topology (By similarity) {ECO:0000250|UniProtKB:P02683, ECO:0000250|UniProtKB:Q62092, ECO:0000269|PubMed:20599942}


Images




Western blot analysis of NSG1 expression in mouse muscle (A), rat muscle (B) whole cell lysates. (Predicted band size: 20 kD; Observed band size: 21 kD)



Immunohistochemical analysis of NSG1 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of NSG1 staining in LS8 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).



Immunohistochemical analysis of NSG1 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.