

Anti-GPR73a Antibody

Rabbit polyclonal antibody to GPR73a

Catalog # AP60619

Product Information

Application	WB, IF/IC
Primary Accession	Q8TCW9
Other Accession	Q9JKL1
Reactivity	Human, Mouse, Rat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	44770

Additional Information

Gene ID	10887
Other Names	GPR73; PKR1; Prokineticin receptor 1; PK-R1; G-protein coupled receptor 73; G-protein coupled receptor ZAQ; GPR73a
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human GPR73a. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000), IF/IC (1/100 - 1/500) IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	PROKR1
Synonyms	GPR73, PKR1
Function	Receptor for prokineticin 1. Exclusively coupled to the G(q) subclass of heteromeric G proteins. Activation leads to mobilization of calcium, stimulation of phosphoinositide turnover and activation of p44/p42 mitogen-activated protein kinase. May play a role during early pregnancy.
Cellular Location	Cell membrane; Multi-pass membrane protein.
Tissue Location	Localizes to glandular epithelium, stroma and vascular endothelial cells of first trimester decidua (at protein level). Up-regulated in first trimester decidua when compared with non- pregnant endometrium. Expressed in the stomach, throughout the small intestine, colon, rectum, thyroid gland,

pituitary gland, salivary gland, adrenal gland, testis, ovary, brain, spleen, prostate and pancreas.

Background

KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human GPR73a. The exact sequence is proprietary.

Images



Western blot analysis of GPR73a expression in BV2 (A), H9C2 (B), HCC827 (C), HEK293T (D) whole cell lysates.



Immunofluorescent analysis of GPR73a staining in Jurkat cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.