

# Anti-IP Receptor Antibody

Rabbit polyclonal antibody to IP Receptor Catalog # AP60621

#### **Product Information**

Application WB, IF/IC
Primary Accession P43119
Reactivity Human, Rat
Host Rabbit
Clonality Polyclonal
Calculated MW 40956

#### **Additional Information**

**Gene ID** 5739

Other Names PRIPR; Prostacyclin receptor; Prostaglandin I2 receptor; PGI receptor; PGI2

receptor; Prostanoid IP receptor

**Target/Specificity** Recognizes endogenous levels of IP Receptor protein.

**Dilution** WB~~WB (1/500 - 1/1000), IF/IC (1/100 - 1/500) IF/IC~~N/A

**Format** Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30%

glycerol, and 0.09% (W/V) sodium azide.

**Storage** Store at -20 °C.Stable for 12 months from date of receipt

#### **Protein Information**

Name PTGIR

Synonyms PRIPR

**Function** Receptor for prostacyclin (prostaglandin I2 or PGI2). The activity of this


receptor is mediated by G(s) proteins which activate adenylate cyclase.

**Cellular Location** Cell membrane; Multi-pass membrane protein.


### **Background**

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human IP Receptor. The exact sequence is proprietary.

## **Images**



Western blot analysis of IP Receptor expression in HEK293T (A), Hela (B), HepG2 (C) whole cell lysates.



Immunofluorescent analysis of IP Receptor staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.