

# Anti-GPR172B Antibody

Rabbit polyclonal antibody to GPR172B

Catalog # AP60634

## Product Information

Application	WB, IF/IC
Primary Accession	<a href="#">Q9NWF4</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	46317

## Additional Information

Gene ID	55065
Other Names	GPR172B; PAR2; RFT1; Solute carrier family 52, riboflavin transporter, member 1; Porcine endogenous retrovirus A receptor 2; PERV-A receptor 2; Protein GPR172B; Riboflavin transporter 1; hRFT1
Target/Specificity	Recognizes endogenous levels of GPR172B protein.
Dilution	WB~~WB (1/500 - 1/1000), IF/IC (1/100 - 1/500) IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

## Protein Information

Name	SLC52A1 ( <a href="#">HGNC:30225</a> )
Synonyms	GPR172B, PAR2, RFT1
Function	Plasma membrane transporter mediating the uptake by cells of the water soluble vitamin B2/riboflavin that plays a key role in biochemical oxidation-reduction reactions of the carbohydrate, lipid, and amino acid metabolism (PubMed: <a href="#">18632736</a> , PubMed: <a href="#">20463145</a> ). Humans are unable to synthesize vitamin B2/riboflavin and must obtain it via intestinal absorption (PubMed: <a href="#">20463145</a> ).
Cellular Location	Cell membrane; Multi-pass membrane protein
Tissue Location	Widely expressed. Highly expressed in the testis, placenta and small intestine. Expressed at lower level in other tissues.


## Background

---


KLH-conjugated synthetic peptide encompassing a sequence within the center region of human GPR172B. The exact sequence is proprietary.

## Images

---



Western blot analysis of GPR172B expression in mouse muscle (A), rat kidney (B) whole cell lysates.



Immunofluorescent analysis of GPR172B staining in HEK293T cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.