

Anti-TNFAIP8L2 Antibody

Rabbit polyclonal antibody to TNFAIP8L2 Catalog # AP60643

Product Information

Application	WB
Primary Accession	<u>Q6P589</u>
Other Accession	<u>Q9D8Y7</u>
Reactivity	Human, Mouse, Rat, Rabbit, Pig
Host	Rabbit
Clonality	Polyclonal
Calculated MW	20556

Additional Information

Gene ID	79626
Other Names	Tumor necrosis factor alpha-induced protein 8-like protein 2; TIPE2; TNF alpha-induced protein 8-like protein 2; TNFAIP8-like protein 2; Inflammation factor protein 20
Target/Specificity	Recognizes endogenous levels of TNFAIP8L2 protein.
Dilution	WB~~WB (1/500 - 1/1000)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	TNFAIP8L2
Function	Acts as a negative regulator of innate and adaptive immunity by maintaining immune homeostasis (PubMed: <u>27043859</u>). Plays a regulatory role in the Toll-like signaling pathway by determining the strength of LPS-induced signaling and gene expression (PubMed: <u>32188758</u>). Inhibits TCR-mediated T-cell activation and negatively regulate T-cell function to prevent hyperresponsiveness (By similarity). Also inhibits autolysosome formation via negatively modulating MTOR activation by interacting with RAC1 and promoting the disassociation of the RAC1-MTOR complex (PubMed: <u>32460619</u>). Plays an essential role in NK-cell biology by acting as a checkpoint and displaying an expression pattern correlating with NK-cell maturation process and by negatively regulating NK-cell maturation and antitumor immunity (By similarity). Mechanistically, suppresses IL-15-triggered mTOR activity in NK-cells (By similarity).

Cellular Location	Cytoplasm. Nucleus. Lysosome
Tissue Location	Expressed in T-cells, B-cells, macrophages, neurons in the brain and brainstem, and stratified squamous epithelia of the esophagus, cervix and skin.

Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human TNFAIP8L2. The exact sequence is proprietary.

Images



Western blot analysis of TNFAIP8L2 expression in mouse liver (A), mouse kidney (B), rat liver (C), rat kidney (D) whole cell lysates.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.