

# Anti-DDX19B Antibody

Rabbit polyclonal antibody to DDX19B Catalog # AP60752

#### **Product Information**

ApplicationWB, IHCPrimary AccessionQ9UMR2

**Reactivity** Human, Mouse

HostRabbitClonalityPolyclonalCalculated MW53927

#### **Additional Information**

**Gene ID** 11269

Other Names DBP5; DDX19; TDBP; ATP-dependent RNA helicase DDX19B; DEAD box RNA

helicase DEAD5; DEAD box protein 19B

**Target/Specificity** KLH-conjugated synthetic peptide encompassing a sequence within the

N-term region of human DDX19B. The exact sequence is proprietary.

**Dilution** WB~~WB (1/500 - 1/1000), IHC (1/100 - 1/200) IHC~~WB (1/500 - 1/1000), IHC

(1/100 - 1/200)

**Format** Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30%

glycerol, and 0.09% (W/V) sodium azide.

**Storage** Store at -20 °C.Stable for 12 months from date of receipt

### **Protein Information**

Name DDX19B

**Synonyms** DBP5, DDX19, TDBP

**Function** ATP-dependent RNA helicase involved in mRNA export from the nucleus

(PubMed: 10428971). Rather than unwinding RNA duplexes, DDX19B functions as a remodeler of ribonucleoprotein particles, whereby proteins bound to nuclear mRNA are dissociated and replaced by cytoplasmic mRNA binding

proteins (PubMed: 10428971).


**Cellular Location** Cytoplasm. Nucleus, nucleoplasm. Note=Associates with the nuclear pore

complex cytoplasmic fibrils


## **Background**

KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human DDX19B. The exact sequence is proprietary.

## **Images**



Western blot analysis of DDX19B expression in LO2 (A), A375 (B), mouse kindey (C) whole cell lysates.



Immunohistochemical analysis of DDX19B staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.