10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-FOXN4 Antibody

Catalog # AP60762

Product Information

Application	WB, IHC
Primary Accession	Q96NZ1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	55215


Additional Information

Gene ID	121643
Other Names	Forkhead box protein N4
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human FOXN4. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000), IHC (1/100 - 1/200) IHC~~WB (1/500 - 1/1000), IHC (1/100 - 1/200)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt


Protein Information

Name	FOXN4
Function	Transcription factor essential for neural and some non-neural tissues development, such as retina and lung respectively. Binds to an 11-bp consensus sequence containing the invariant tetranucleotide 5'- ACGC-3'. During development of the central nervous system, is required to specify the amacrine and horizontal cell fates from multipotent retinal progenitors while suppressing the alternative photoreceptor cell fates through activating DLL4-NOTCH signaling. Also acts synergistically with ASCL1/MASH1 to activate DLL4-NOTCH signaling and drive commitment of p2 progenitors to the V2b interneuron fates during spinal cord neurogenesis. In development of non-neural tissues, plays an essential role in the specification of the atrioventricular canal and is indirectly required for patterning the distal airway during lung development (By similarity).
Cellular Location	Nucleus {ECO:0000255|PROSITE-ProRule:PRU00089}.


Images



Western blot analysis of FOXN4 expression in H1792 (A) whole cell lysates. (Predicted band size: 55 kD; Observed band size: 55 kD)



Immunohistochemical analysis of FOXN4 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunohistochemical analysis of FOXN4 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.