

Anti-GIP Receptor Antibody

Rabbit polyclonal antibody to GIP Receptor

Catalog # AP60804

Product Information

Application	WB, IF/IC, IHC
Primary Accession	P48546
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	53157

Additional Information


Gene ID	2696
Other Names	Gastric inhibitory polypeptide receptor; GIP-R; Glucose-dependent insulinotropic polypeptide receptor
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human GIP Receptor. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500) IF/IC~~N/A IHC~~WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	GIPR
Function	This is a receptor for GIP. The activity of this receptor is mediated by G proteins which activate adenylyl cyclase.
Cellular Location	Cell membrane; Multi-pass membrane protein

Background


KLH-conjugated synthetic peptide encompassing a sequence within the center region of human GIP Receptor. The exact sequence is proprietary.



Western blot analysis of GIP Receptor expression in HepG2 (A) whole cell lysates.



Immunohistochemical analysis of GIP Receptor staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of GIP Receptor staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.