

Anti-CD107b Antibody

Rabbit polyclonal antibody to CD107b
Catalog # AP60811

Product Information

Application	WB
Primary Accession	P13473
Other Accession	P17047
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	44961

Additional Information

Gene ID	3920
Other Names	Lysosome-associated membrane glycoprotein 2; LAMP-2; Lysosome-associated membrane protein 2; CD107 antigen-like family member B; CD107b
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human CD107b. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/2000)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C. Stable for 12 months from date of receipt

Protein Information

Name	LAMP2
Function	Lysosomal membrane glycoprotein which plays an important role in lysosome biogenesis, lysosomal pH regulation and autophagy (PubMed: 11082038 , PubMed: 18644871 , PubMed: 24880125 , PubMed: 27628032 , PubMed: 36586411 , PubMed: 37390818 , PubMed: 8662539). Acts as an important regulator of lysosomal lumen pH regulation by acting as a direct inhibitor of the proton channel TMEM175, facilitating lysosomal acidification for optimal hydrolase activity (PubMed: 37390818). Plays an important role in chaperone-mediated autophagy, a process that mediates lysosomal degradation of proteins in response to various stresses and as part of the normal turnover of proteins with a long biological half-life (PubMed: 11082038 , PubMed: 18644871 , PubMed: 24880125 , PubMed: 27628032 , PubMed: 36586411 ,

PubMed:[8662539](#)). Functions by binding target proteins, such as GAPDH, GPX4, NLRP3 and MLLT11, and targeting them for lysosomal degradation (PubMed:[11082038](#), PubMed:[18644871](#), PubMed:[24880125](#), PubMed:[36586411](#), PubMed:[8662539](#)). In the chaperone-mediated autophagy, acts downstream of chaperones, such as HSPA8/HSC70, which recognize and bind substrate proteins and mediate their recruitment to lysosomes, where target proteins bind LAMP2 (PubMed:[36586411](#)). Plays a role in lysosomal protein degradation in response to starvation (By similarity). Required for the fusion of autophagosomes with lysosomes during autophagy (PubMed:[27628032](#)). Cells that lack LAMP2 express normal levels of VAMP8, but fail to accumulate STX17 on autophagosomes, which is the most likely explanation for the lack of fusion between autophagosomes and lysosomes (PubMed:[27628032](#)). Required for normal degradation of the contents of autophagosomes (PubMed:[27628032](#)). Required for efficient MHC class II-mediated presentation of exogenous antigens via its function in lysosomal protein degradation; antigenic peptides generated by proteases in the endosomal/lysosomal compartment are captured by nascent MHC II subunits (PubMed:[15894275](#), PubMed:[20518820](#)). Is not required for efficient MHC class II-mediated presentation of endogenous antigens (PubMed:[20518820](#)).

Cellular Location

Lysosome membrane {ECO:0000255 | PROSITE- ProRule:PRU00740, ECO:0000269 | PubMed:11082038, ECO:0000269 | PubMed:17897319, ECO:0000269 | PubMed:18644871, ECO:0000269 | PubMed:2912382}; Single-pass type I membrane protein {ECO:0000255 | PROSITE-ProRule:PRU00740, ECO:0000269 | PubMed:17897319} Endosome membrane; Single-pass type I membrane protein {ECO:0000255 | PROSITE-ProRule:PRU00740, ECO:0000269 | PubMed:17897319}. Cell membrane; Single-pass type I membrane protein {ECO:0000255 | PROSITE-ProRule:PRU00740, ECO:0000269 | PubMed:17897319}. Cytoplasmic vesicle, autophagosome membrane {ECO:0000250 | UniProtKB:P17047}. Note=This protein shuttles between lysosomes, endosomes, and the plasma membrane

Tissue Location


Isoform LAMP-2A is highly expressed in placenta, lung and liver, less in kidney and pancreas, low in brain and skeletal muscle (PubMed:26856698, PubMed:7488019). Isoform LAMP-2B is detected in spleen, thymus, prostate, testis, small intestine, colon, skeletal muscle, brain, placenta, lung, kidney, ovary and pancreas and liver (PubMed:26856698, PubMed:7488019). Isoform LAMP-2C is detected in small intestine, colon, heart, brain, skeletal muscle, and at lower levels in kidney and placenta (PubMed:26856698).

Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human CD107b. The exact sequence is proprietary.

Images

Western blot analysis of CD107b expression in A375 (A), DLD (B), mouse lung (C), mouse liver (D), rat lung (E), rat liver (F) whole cell lysates.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.