

# Anti-AT1 Antibody

Rabbit polyclonal antibody to AT1

Catalog # AP60812

## Product Information

Application	WB, IF/IC, IHC
Primary Accession	<a href="#">P30556</a>
Reactivity	Human, Mouse, Rat, Pig, Bovine, SARS
Host	Rabbit
Clonality	Polyclonal
Calculated MW	41061

## Additional Information

Gene ID	185
Other Names	AGTR1A; AGTR1B; AT2R1; AT2R1B; Type-1 angiotensin II receptor; AT1AR; AT1BR; Angiotensin II type-1 receptor; AT1
Target/Specificity	Recognizes endogenous levels of AT1 protein.
Dilution	WB~~WB (1/500 - 1/2000), IHC (1/50 - 1/200), IF/IC (1/50 - 1/100) IF/IC~~N/A IHC~~WB (1/500 - 1/2000), IHC (1/50 - 1/200), IF/IC (1/50 - 1/100)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt


## Protein Information

Name	AGTR1 ( <a href="#">HGNC:336</a> )
Function	Receptor for angiotensin II, a vasoconstricting peptide, which acts as a key regulator of blood pressure and sodium retention by the kidney (PubMed: <a href="#">15611106</a> , PubMed: <a href="#">1567413</a> , PubMed: <a href="#">25913193</a> , PubMed: <a href="#">26420482</a> , PubMed: <a href="#">30639100</a> , PubMed: <a href="#">32079768</a> , PubMed: <a href="#">8987975</a> ). The activated receptor in turn couples to G-alpha proteins G(q) (GNAQ, GNA11, GNA14 or GNA15) and thus activates phospholipase C and increases the cytosolic Ca(2+) concentrations, which in turn triggers cellular responses such as stimulation of protein kinase C (PubMed: <a href="#">15611106</a> ).
Cellular Location	Cell membrane; Multi-pass membrane protein
Tissue Location	Liver, lung, adrenal and adrenocortical adenomas.


## Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human AT1. The exact sequence is proprietary.


## Images



Western blot analysis of AT1 expression in C6 (A), PC3 (B), U87MG (C), HCC827 (D) whole cell lysates.



Immunohistochemical analysis of AT1 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of AT1 staining in K562 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.