

# Anti-Apolipoprotein A5 Antibody

Rabbit polyclonal antibody to Apolipoprotein A5 Catalog # AP60839

# **Product Information**

| Application       | WB                        |
|-------------------|---------------------------|
| Primary Accession | <u>Q6Q788</u>             |
| Other Accession   | <u>Q8C7G5</u>             |
| Reactivity        | Human, Mouse, Rat, Monkey |
| Host              | Rabbit                    |
| Clonality         | Polyclonal                |
| Calculated MW     | 41213                     |
|                   | 5                         |

## **Additional Information**

| Gene ID            | 116519  |
|--------------------|---|
| Other Names        | RAP3; Apolipoprotein A-V; Apo-AV; ApoA-V; Apolipoprotein A5;<br>Regeneration-associated protein 3               |
| Target/Specificity | Recognizes endogenous levels of Apolipoprotein A5 protein.  |
| Dilution           | WB~~WB (1/500 - 1/2000)   |
| Format             | Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide. |
| Storage            | Store at -20 °C.Stable for 12 months from date of receipt   |

### **Protein Information**

| Name              | APOA5   |
|-------------------|---|
| Synonyms          | RAP3  |
| Function          | Minor apolipoprotein mainly associated with HDL and to a lesser extent with VLDL. May also be associated with chylomicrons. Important determinant of plasma triglyceride (TG) levels by both being a potent stimulator of apo-CII lipoprotein lipase (LPL) TG hydrolysis and an inhibitor of the hepatic VLDL-TG production rate (without affecting the VLDL-apoB production rate) (By similarity). Activates poorly lecithin:cholesterol acyltransferase (LCAT) and does not enhance efflux of cholesterol from macrophages. Binds heparin (PubMed: <u>17326667</u> ). |
| Cellular Location | Secreted. Early endosome. Late endosome. Golgi apparatus, trans-Golgi<br>network. Note=In the presence of SORL1, internalized to early endosomes,<br>sorted in a retrograde fashion to late endosomes, from which a portion is  |

sent to lysosomes and degradation, another portion is sorted to the trans-Golgi network

**Tissue Location** 

Liver and plasma.

#### Background

KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human Apolipoprotein A5. The exact sequence is proprietary.

#### Images



Western blot analysis of Apolipoprotein A5 expression in HEK293T (A), H446 (B), Jurkat (C), mouse liver (D), mouse heart (E), rat liver (F), rat heart (G) whole cell lysates.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.