

Anti-IRS2 Antibody

Rabbit polyclonal antibody to IRS2

Catalog # AP60877

Product Information

Application	WB
Primary Accession	Q9Y4H2
Other Accession	P81122
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	137334

Additional Information

Gene ID	8660
Other Names	Insulin receptor substrate 2; IRS-2
Target/Specificity	Recognizes endogenous levels of IRS2 protein.
Dilution	WB~~WB (1/500 - 1/1000)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	IRS2
Function	Signaling adapter protein that participates in the signal transduction from two prominent receptor tyrosine kinases, insulin receptor/INSR and insulin-like growth factor I receptor/IGF1R (PubMed: 25879670). Plays therefore an important role in development, growth, glucose homeostasis as well as lipid metabolism (PubMed: 24616100). Upon phosphorylation by the insulin receptor, functions as a signaling scaffold that propagates insulin action through binding to SH2 domain-containing proteins including the p85 regulatory subunit of PI3K, NCK1, NCK2, GRB2 or SHP2 (PubMed: 15316008 , PubMed: 19109239). Recruitment of GRB2 leads to the activation of the guanine nucleotide exchange factor SOS1 which in turn triggers the Ras/Raf/MEK/MAPK signaling cascade (By similarity). Activation of the PI3K/AKT pathway is responsible for most of insulin metabolic effects in the cell, and the Ras/Raf/MEK/MAPK is involved in the regulation of gene expression and in cooperation with the PI3K pathway regulates cell growth and differentiation. Acts a positive regulator of the Wnt/beta- catenin

signaling pathway through suppression of DVL2 autophagy- mediated degradation leading to cell proliferation (PubMed:[24616100](#)). Plays a role in cell cycle progression by promoting a robust spindle assembly checkpoint (SAC) during M-phase (PubMed:[32554797](#)). In macrophages, IL4-induced tyrosine phosphorylation of IRS2 leads to the recruitment and activation of phosphoinositide 3-kinase (PI3K) (PubMed:[19109239](#)).


Cellular Location

Cytoplasm, cytosol {ECO:0000250 | UniProtKB:P81122}

Background

KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human IRS2. The exact sequence is proprietary.

Images



Western blot analysis of IRS2 expression in HEK293T (A) whole cell lysates.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.