

Anti-TRIM59 Antibody

Rabbit polyclonal antibody to TRIM59

Catalog # AP60887

Product Information

Application	WB, IF/IC, IHC
Primary Accession	Q8IWR1
Other Accession	Q922Y2
Reactivity	Human, Mouse, Rat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	47114

Additional Information

Gene ID	286827
Other Names	RNF104; TRIM57; TSBF1; Tripartite motif-containing protein 59; RING finger protein 104; Tumor suppressor TSBF-1
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human TRIM59. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000), IHC (1/50 - 1/100), IF/IC (1/100 - 1/500) IF/IC~~N/A IHC~~WB (1/500 - 1/1000), IHC (1/50 - 1/100), IF/IC (1/100 - 1/500)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	TRIM59
Synonyms	RNF104, TRIM57, TSBF1
Function	E3 ubiquitin ligase involved in different processes such as development and immune response (PubMed: 22588174 , PubMed: 30231667). Serves as a negative regulator for innate immune signaling pathways by suppressing RLR-induced activation of IRF3/7 and NF-kappa-B via interaction with adapter ECSIT (PubMed: 22588174). Regulates autophagy through modulating both the transcription and the ubiquitination of BECN1 (PubMed: 30231667). On the one hand, regulates the transcription of BECN1 through negatively modulating the NF-kappa-B pathway. On the other hand, regulates TRAF6-mediated 'Lys-63'-linked ubiquitination of BECN1, thus affecting the formation of the BECN1-PIK3C3 complex. In addition, mediates 'Lys-48'-linked

ubiquitination of TRAF6 and thereby promotes TRAF6 proteasomal degradation (PubMed:[30231667](#)). Also acts as a critical regulator for early embryo development from blastocyst stage to gastrula through modulating F-actin assembly and WASH1 'Lys-63'- linked ubiquitination (By similarity).


Cellular Location

Endoplasmic reticulum membrane; Single-pass membrane protein


Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human TRIM59. The exact sequence is proprietary.


Images



Western blot analysis of TRIM59 expression in Jurkat (A), HEK293T (B), rat brain (C), mouse brain (D) whole cell lysates.



Immunohistochemical analysis of TRIM59 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of TRIM59 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with Alexa Fluor 647-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.