

Anti-ZAK (pS165) Antibody

Rabbit polyclonal antibody to ZAK (pS165) Catalog # AP60932

Product Information

Application WB
Primary Accession Q9NYL2
Reactivity Human
Host Rabbit
Clonality Polyclonal
Calculated MW 91155

Additional Information

Gene ID 51776

Other Names MLTK; Mitogen-activated protein kinase kinase kinase MLT; Human cervical

cancer suppressor gene 4 protein; HCCS-4; Leucine zipper- and sterile alpha motif-containing kinase; MLK-like mitogen-activated protein triple kinase; Mixed lineage kinase-related kinase; MLK-related kinase; MRK; Sterile alpha

motif- and leucine zipper-containing kinase AZK

Target/Specificity KLH-conjugated synthetic peptide encompassing a sequence within the center

region of human ZAK. The exact sequence is proprietary.

Dilution WB~~WB (1/500 - 1/1000)

Format Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30%

glycerol, and 0.09% (W/V) sodium azide.

Storage Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name MAP3K20 (<u>HGNC:17797</u>)

Function Stress-activated component of a protein kinase signal transduction cascade

that promotes programmed cell death in response to various stress, such as ribosomal stress, osmotic shock and ionizing radiation (PubMed: 10924358,

PubMed:<u>11836244</u>, PubMed:<u>12220515</u>, PubMed:<u>14521931</u>, PubMed:<u>15350844</u>, PubMed:<u>15737997</u>, PubMed:<u>18331592</u>, PubMed:<u>20559024</u>, PubMed:<u>26999302</u>, PubMed:<u>32289254</u>,

PubMed:32610081, PubMed:35857590). Acts by catalyzing phosphorylation of

MAP kinase kinases, leading to activation of the JNK (MAPK8/JNK1,

MAPK9/JNK2 and/or MAPK10/JNK3) and MAP kinase p38 (MAPK11, MAPK12, MAPK13 and/or MAPK14) pathways (PubMed: 11042189, PubMed: 11836244,

PubMed: 12220515, PubMed: 14521931, PubMed: 15172994,

PubMed:<u>15737997</u>, PubMed:<u>32289254</u>, PubMed:<u>32610081</u>, PubMed:<u>35857590</u>). Activates JNK through phosphorylation of MAP2K4/MKK4 and MAP2K7/MKK7, and MAP kinase p38 gamma (MAPK12) via phosphorylation of MAP2K3/MKK3 and MAP2K6/MKK6 (PubMed:<u>11836244</u>, PubMed:<u>12220515</u>). Involved in stress associated with adrenergic stimulation: contributes to cardiac decompensation during periods of acute cardiac stress (PubMed:<u>15350844</u>, PubMed:<u>21224381</u>, PubMed:<u>27859413</u>). May be involved in regulation of S and G2 cell cycle checkpoint by mediating phosphorylation of CHEK2 (PubMed:<u>15342622</u>).

Cellular Location

Cytoplasm. Nucleus. Note=Translocates to the nucleus upon ultraviolet B

irradiation.


Tissue Location

Ubiquitously expressed. Isoform ZAKbeta is the predominant form in all tissues examined, except for liver, in which isoform ZAKalpha is more highly expressed

Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human ZAK. The exact sequence is proprietary.

Images



Western blot analysis of ZAK (pS165) expression in SGC7901 (A), HEK293T (B), HCT116 (C) whole cell lysates.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.