10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-CDC14A Antibody

Catalog # AP61024

Product Information

Application	WB, IF/IC
Primary Accession	Q9UNH5
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	66574

Additional Information


Gene ID	8556
Other Names	Dual specificity protein phosphatase CDC14A; CDC14 cell division cycle 14 homolog A
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human CDC14A. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	CDC14A
Function	Dual-specificity phosphatase. Required for centrosome separation and productive cytokinesis during cell division. Dephosphorylates SIRT2 around early anaphase. May dephosphorylate the APC subunit FZR1/CDH1, thereby promoting APC-FZR1 dependent degradation of mitotic cyclins and subsequent exit from mitosis. Required for normal hearing (PubMed:29293958).
Cellular Location	Nucleus. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle pole. Cytoplasm, cytoskeleton, spindle. Cell projection, kinocilium {ECO:0000250|UniProtKB:Q6GQT0}. Cell projection, stereocilium {ECO:0000250|UniProtKB:Q6GQT0}. Note=Centrosomal during interphase, released into the cytoplasm at the onset of mitosis. Subsequently localizes to the mitotic spindle pole and at the central spindle (PubMed:11901424, PubMed:12134069, PubMed:15263015). Present along both the transient kinocilia of developing cochlear hair cells

and the persistent kinocilia of vestibular hair cells (By similarity) {ECO:0000250|UniProtKB:Q6GQT0, ECO:0000269|PubMed:11901424, ECO:0000269|PubMed:12134069, ECO:0000269|PubMed:15263015}

Images



Western blot analysis of CDC14A expression in C6 (A), MEF (B), K562 (C), HEK293T (D) whole cell lysates. (Predicted band size: 66 kD; Observed band size: 67 kD)



Immunofluorescent analysis of CDC14A staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.