

Anti-Progesterone Receptor (pS400) Antibody

Rabbit polyclonal antibody to Progesterone Receptor (pS400)

Catalog # AP61145

Product Information

Application	WB, IF/IC
Primary Accession	P06401
Other Accession	Q00175
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	98981

Additional Information

Gene ID	5241
Other Names	NR3C3; Progesterone receptor; PR; Nuclear receptor subfamily 3 group C member 3
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Progesterone Receptor (pS400). The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000), IF/IC (1/100 - 1/500) IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	PGR
Synonyms	NR3C3
Function	The steroid hormones and their receptors are involved in the regulation of eukaryotic gene expression and affect cellular proliferation and differentiation in target tissues. Depending on the isoform, progesterone receptor functions as a transcriptional activator or repressor.
Cellular Location	Nucleus. Cytoplasm. Note=Nucleoplasmic shuttling is both hormone- and cell cycle-dependent. On hormone stimulation, retained in the cytoplasm in the G(1) and G(2)/M phases [Isoform 4]: Mitochondrion outer membrane In reproductive tissues the expression of isoform A and isoform B varies as a


Tissue Location

consequence of developmental and hormonal status. Isoform A and isoform B are expressed in comparable levels in uterine glandular epithelium during the proliferative phase of the menstrual cycle. Expression of isoform B but not of isoform A persists in the glands during mid-secretory phase. In the stroma, isoform A is the predominant form throughout the cycle. Heterogeneous isoform expression between the glands of the endometrium basalis and functionalis is implying region-specific responses to hormonal stimuli


Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Progesterone Receptor (pS400). The exact sequence is proprietary.

Images



Western blot analysis of Progesterone Receptor (pS400) expression in mouse testis (A), rat ovary (B), A2780 (C), HeLa (D) whole cell lysates.



Immunofluorescent analysis of Progesterone Receptor (pS400) staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a Alexa Fluor 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.