10320 Camino Santa Fe, Suite G San Diego, CA 92121 Tel: 858.875.1900 Fax: 858.875.1999



Anti-CHRM2 Antibody

Rabbit polyclonal antibody to CHRM2 Catalog # AP61231

Product Information

ApplicationWB, IF/ICPrimary AccessionP08172Other AccessionO9ERZ4

Reactivity Human, Mouse, Rat, Pig, Bovine

HostRabbitClonalityPolyclonalCalculated MW51715

Additional Information

Gene ID 1129

Other Names Muscarinic acetylcholine receptor M2

Target/Specificity Recognizes endogenous levels of CHRM2 protein.

Dilution WB~~WB (1/500 - 1/1000), IF/IC (1/100 - 1/500) IF/IC~~N/A

Format Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30%

glycerol, and 0.09% (W/V) sodium azide.

Storage Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name CHRM2

Function The muscarinic acetylcholine receptor mediates various cellular responses,

including inhibition of adenylate cyclase, breakdown of phosphoinositides and modulation of potassium channels through the action of G proteins. Primary transducing effect is adenylate cyclase inhibition. Signaling promotes phospholipase C activity, leading to the release of inositol trisphosphate (IP3);

this then triggers calcium ion release into the cytosol.


Cellular Location Cell membrane; Multi-pass membrane protein. Postsynaptic cell membrane;

Multi-pass membrane protein. Note=Phosphorylation in response to agonist binding promotes receptor internalization {ECO:0000250|UniProtKB:P06199}

Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human CHRM2. The exact sequence is proprietary.

Images



Western blot analysis of CHRM2 expression in HEK293T (A), mouse lung (B), mouse kidney (C), rat kidney (D) whole cell lysates.



Immunofluorescent analysis of CHRM2 staining in LOVO cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a Alexa Fluor 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.