

# Anti-IL-11 Antibody

Rabbit polyclonal antibody to IL-11

Catalog # AP61233

## Product Information

---

Application	WB, IHC
Primary Accession	<a href="#">P20809</a>
Other Accession	<a href="#">P47873</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	21429

## Additional Information

---

Gene ID	3589
Other Names	Interleukin-11; IL-11; Adipogenesis inhibitory factor; AGIF; Oprelvekin
Target/Specificity	Recognizes endogenous levels of IL-11 protein.
Dilution	WB~~WB (1/500 - 1/1000), IHC (1/50 - 1/200) IHC~~WB (1/500 - 1/1000), IHC (1/50 - 1/200)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

## Protein Information

---

Name	IL11 ( <a href="#">HGNC:5966</a> )
Function	Cytokine that stimulates the proliferation of hematopoietic stem cells and megakaryocyte progenitor cells and induces megakaryocyte maturation resulting in increased platelet production (PubMed: <a href="#">2145578</a> ). Also promotes the proliferation of hepatocytes in response to liver damage. Binding to its receptor formed by IL6ST and IL11RA activates a signaling cascade that promotes cell proliferation (PubMed: <a href="#">12919066</a> ). Signaling leads to the activation of intracellular protein kinases and the phosphorylation of STAT3. The interaction with the membrane-bound IL11RA and IL6ST stimulates 'classic signaling', whereas the binding of IL11 and soluble IL11RA to IL6ST stimulates 'trans-signaling' (PubMed: <a href="#">30279168</a> ).
Cellular Location	Secreted


## Background

---


KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human IL-11. The exact sequence is proprietary.

## Images

---



Western blot analysis of IL-11 expression in 3T3L1 (A), HeLa (B), PC12 (C) whole cell lysates.



Immunohistochemical analysis of IL-11 staining in human pancreas formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.