

Anti-NEFH Antibody

Rabbit polyclonal antibody to NEFH

Catalog # AP61237

Product Information

| | |
|-------------------|------------------------|
| Application | WB, IC, IHC |
| Primary Accession | P12036 |
| Other Accession | P19246 |
| Reactivity | Human, Mouse, Rat |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 111838 |

Additional Information

| | |
|--------------------|---|
| Gene ID | 4744 |
| Other Names | KIAA0845; NFH; Neurofilament heavy polypeptide; NF-H; 200 kDa neurofilament protein; Neurofilament triplet H protein |
| Target/Specificity | Recognizes endogenous levels of NEFH protein. |
| Dilution | WB~~WB (1/500 - 1/1000), IHC (1/50 - 1/200), IC (1/100 - 1/500) IC~~N/A IHC~~WB (1/500 - 1/1000), IHC (1/50 - 1/200), IC (1/100 - 1/500) |
| Format | Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide. |
| Storage | Store at -20 °C.Stable for 12 months from date of receipt |


Protein Information

| | |
|-------------------|---|
| Name | NEFH |
| Synonyms | KIAA0845, NFH |
| Function | Neurofilaments usually contain three intermediate filament proteins: NEFL, NEFM, and NEFH which are involved in the maintenance of neuronal caliber. NEFH has an important function in mature axons that is not subserved by the two smaller NF proteins. May additionally cooperate with the neuronal intermediate filament proteins PRPH and INA to form neuronal filamentous networks (By similarity). |
| Cellular Location | Cytoplasm, cytoskeleton. Cell projection, axon {ECO:0000250 UniProtKB:P19246} |


Background

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human NEFH. The exact sequence is proprietary.


Images



Western blot analysis of NEFH expression in C6 (A), CT26 (B), HepG2 (C), A2780 (D) whole cell lysates.



Immunohistochemical analysis of NEFH staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunocytochemistry analysis of NEFH staining in NIH3T3 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a HRP-conjugated secondary antibody in PBS at room temperature. DAB was used as the chromogen.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.