

# Anti-MLL1 Antibody

Rabbit polyclonal antibody to MLL1 Catalog # AP61295

#### **Product Information**

ApplicationWBPrimary AccessionQ03164Other AccessionP55200

**Reactivity** Human, Mouse, Rat, Monkey, Pig, Bovine, Drosophila

Host Rabbit
Clonality Polyclonal
Calculated MW 431764

### **Additional Information**

**Gene ID** 4297

Other Names ALL1; CXXC7; HRX; HTRX; MLL; MLL1; TRX1; Histone-lysine

N-methyltransferase 2A; Lysine N-methyltransferase 2A; ALL-1; CXXC-type zinc

finger protein 7; Myeloid/lymphoid or mixed-lineage leukemia;

Myeloid/lymphoid or mixed-lineage leukemia protein 1; Trithorax-like protein;

Zinc finger protein HRX

**Target/Specificity** KLH-conjugated synthetic peptide encompassing a sequence within the

C-term region of human MLL1. The exact sequence is proprietary.

**Dilution** WB~~WB (1/500 - 1/1000)

**Format** Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30%

glycerol, and 0.09% (W/V) sodium azide.

**Storage** Store at -20 °C.Stable for 12 months from date of receipt

#### **Protein Information**

Name KMT2A

Synonyms ALL1, CXXC7, HRX, HTRX, MLL, MLL1, TRX1

**Function** Histone methyltransferase that plays an essential role in early development

and hematopoiesis (PubMed:<u>12453419</u>, PubMed:<u>15960975</u>, PubMed:<u>19187761</u>, PubMed:<u>19556245</u>, PubMed:<u>20677832</u>,

PubMed:<u>21220120</u>, PubMed:<u>26886794</u>). Catalytic subunit of the MLL1/MLL complex, a multiprotein complex that mediates both methylation of 'Lys-4' of histone H3 (H3K4me) complex and acetylation of 'Lys-16' of histone H4 (H4K16ac) (PubMed:<u>12453419</u>, PubMed:<u>15960975</u>, PubMed:<u>19187761</u>,

PubMed: 19556245, PubMed: 20677832, PubMed: 21220120,

PubMed: 24235145, PubMed: 26886794). Catalyzes methyl group transfer from S-adenosyl-L- methionine to the epsilon-amino group of 'Lys-4' of histone H3 (H3K4) via a non-processive mechanism. Part of chromatin remodeling machinery predominantly forms H3K4me1 and H3K4me2 methylation marks at active chromatin sites where transcription and DNA repair take place (PubMed:12453419, PubMed:15960975, PubMed:19187761, PubMed: 19556245, PubMed: 20677832, PubMed: 21220120, PubMed: <u>25561738</u>, PubMed: <u>26886794</u>). Has weak methyltransferase activity by itself, and requires other component of the MLL1/MLL complex to obtain full methyltransferase activity (PubMed:19187761, PubMed:26886794). Has no activity toward histone H3 phosphorylated on 'Thr-3', less activity toward H3 dimethylated on 'Arg-8' or 'Lys-9', while it has higher activity toward H3 acetylated on 'Lys-9' (PubMed: 19187761). Binds to unmethylated CpG elements in the promoter of target genes and helps maintain them in the nonmethylated state (PubMed: 20010842). Required for transcriptional activation of HOXA9 (PubMed: 12453419, PubMed: 20010842, PubMed: 20677832). Promotes PPP1R15A-induced apoptosis (PubMed: 10490642). Plays a critical role in the control of circadian gene expression and is essential for the transcriptional activation mediated by the CLOCK-BMAL1 heterodimer (By similarity). Establishes a permissive chromatin state for circadian transcription by mediating a rhythmic methylation of 'Lys-4' of histone H3 (H3K4me) and this histone modification directs the circadian acetylation at H3K9 and H3K14 allowing the recruitment of CLOCK-BMAL1 to chromatin (By similarity). Also has auto-methylation activity on Cys-3882 in absence of histone H3 substrate (PubMed: 24235145).

**Cellular Location** 

Nucleus [MLL cleavage product C180]: Nucleus. Note=Localizes to a diffuse nuclear pattern when not associated with MLL cleavage product N320


**Tissue Location** 

Heart, lung, brain and T- and B-lymphocytes.

## **Background**

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human MLL1. The exact sequence is proprietary.

## **Images**



Western blot analysis of MLL1 expression in K562 (A), A375 (B), U87MG (C), mouse brain (D), rat spleen (E) whole cell lysates.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.