10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-Alpha-2B Adrenergic Receptor Antibody

Catalog # AP61305

Product Information

Application	WB, IHC, IF/IC
Primary Accession	P18089
Reactivity	Human, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	49954

Additional Information

Gene ID	151
Other Names	ADRA2L1; ADRA2RL1; Alpha-2B adrenergic receptor; Alpha-2 adrenergic receptor subtype C2; Alpha-2B adrenoreceptor; Alpha-2B adrenoceptor; Alpha-2BAR
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Alpha-2B Adrenergic Receptor. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000), IF/IC (1/100 - 1/500) IHC~~1:100~500 IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	ADRA2B (HGNC:282)
Synonyms	ADRA2L1, ADRA2RL1
Function	Alpha-2 adrenergic receptors are G protein-coupled receptors for catecholamines that activate G(i/o) protein pathway, thereby promoting adenylyl cyclase inhibition, ERK1/2 stimulation, and voltage- gated calcium channels suppression (PubMed:11056163, PubMed:2164221, PubMed:2172775, PubMed:23105096, PubMed:26811329). Control a variety of physiological processes, such as regulation of blood pressure, lipolysis and insulin release (PubMed:2164221). The rank order of potency for agonists of ADRA2B is clonidine > norepinephrine > epinephrine = oxymetazoline > dopamine > p-tyramine = phenylephrine > serotonin > p-synephrine / p-octopamine. For antagonists, the rank order is yohimbine > chlorpromazine > phentolamine > mianserine > spiperone > prazosin > alprenolol >

propanolol > pindolol (PubMed:2164221, PubMed:2172775).


Cellular Location

Cell membrane; Multi-pass membrane protein Note=Interaction with RAB26, GGA1, GGA2 and GGA3 mediates transport from the Golgi to the cell membrane.


Images



Western blot analysis of Alpha-2B Adrenergic Receptor expression in Hela (A) whole cell lysates. (Predicted band size: 49 kD; Observed band size: 62 kD)



Immunohistochemical analysis of Alpha-2B Adrenergic Receptor staining in human kidney cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of Alpha-2B Adrenergic Receptor staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a Alexa Fluor 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.