10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# Anti-Granulin Antibody

Catalog # AP61369

## Product Information

| | |
|---|---|
| **Application** | WB, IF/IC |
| **Primary Accession** | P28799 |
| **Reactivity** | Human, Rat |
| **Host** | Rabbit |
| **Clonality** | Polyclonal |
| **Calculated MW** | 63544 |

## Additional Information

| | |
|---|---|
| **Gene ID** | 2896 |
| **Other Names** | Granulins; Proepithelin; PEPI |
| **Target/Specificity** | KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Granulin. The exact sequence is proprietary. |
| **Dilution** | WB~~WB (1/500 - 1/1000) IF/IC~~N/A |
| **Format** | Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide. |
| **Storage** | Store at -20 °C.Stable for 12 months from date of receipt |

## Protein Information


| | |
|---|---|
| **Name** | GRN ( HGNC:4601) |
| **Function** | Secreted protein that acts as a key regulator of lysosomal function and as a growth factor involved in inflammation, wound healing and cell proliferation (PubMed:12526812, PubMed:18378771, PubMed:28073925, PubMed:28453791, PubMed:28541286). Regulates protein trafficking to lysosomes, and also the activity of lysosomal enzymes (PubMed:28453791, PubMed:28541286). Also facilitates the acidification of lysosomes, causing degradation of mature CTSD by CTSB (PubMed:28073925). In addition, functions as a wound-related growth factor that acts directly on dermal fibroblasts and endothelial cells to promote division, migration and the formation of capillary-like tubule structures (By similarity). Also promotes epithelial cell proliferation by blocking TNF-mediated neutrophil activation preventing release of oxidants and proteases (PubMed:12526812). Moreover, modulates inflammation in neurons by preserving neurons survival, axonal outgrowth and neuronal integrity (PubMed:18378771). |
| **Cellular Location** | Secreted. Lysosome Note=Endocytosed by SORT1 and delivred to lysosomes |

(PubMed:21092856, PubMed:28073925). Targeted to lysosome by PSAP via M6PR and LRP1, in both biosynthetic and endocytic pathways (PubMed:26370502, PubMed:28073925). Co-localized with GBA1 in the intracellular trafficking compartments until to lysosome (By similarity) {ECO:0000250|UniProtKB:P28798, ECO:0000269|PubMed:21092856, ECO:0000269|PubMed:26370502, ECO:0000269|PubMed:28073925}


**Tissue Location**

In myelogenous leukemic cell lines of promonocytic, promyelocytic, and proerythroid lineage, in fibroblasts, and very strongly in epithelial cell lines. Present in inflammatory cells and bone marrow. Highest levels in kidney

## Images




Western blot analysis of Granulin expression in rat kidney (A), HEK293T (B), Jurkat (C), Beas2B (D) whole cell lysates. (Predicted band size: 63 kD; Observed band size: 50 kD)




Immunofluorescent analysis of Granulin staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.