

Anti-NMDAR1 (Phospho-S890) Antibody

Catalog # AP61377

Product Information

Application	WB, IHC, IF/IC
Primary Accession	Q05586
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	105373

Additional Information

Gene ID	2902
Other Names	NMDAR1; Glutamate receptor ionotropic NMDA 1; GluN1; Glutamate [NMDA] receptor subunit zeta-1; N-methyl-D-aspartate receptor subunit NR1; NMD-R1
Target/Specificity	KLH-conjugated synthetic phosphopeptide corresponding to residues surrounding S890 of human NMDAR1 protein. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000), IHC (1/50 - 1/200), IF/IC (1/100 - 1/500) IHC~~WB (1/500 - 1/1000), IHC (1/50 - 1/200), IF/IC (1/100 - 1/500) IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	GRIN1 (HGNC:4584)
Function	Component of N-methyl-D-aspartate (NMDA) receptors (NMDARs) that function as heterotetrameric, ligand-gated cation channels with high calcium permeability and voltage-dependent block by Mg(2+) (PubMed: 21376300 , PubMed: 26875626 , PubMed: 26919761 , PubMed: 28126851 , PubMed: 28228639 , PubMed: 36959261 , PubMed: 7679115 , PubMed: 7681588 , PubMed: 7685113). NMDARs participate in synaptic plasticity for learning and memory formation by contributing to the long-term potentiation (LTP) (PubMed: 26875626). Channel activation requires binding of the neurotransmitter L-glutamate to the GluN2 subunit, glycine or D-serine binding to the GluN1 subunit, plus membrane depolarization to eliminate channel inhibition by Mg(2+) (PubMed: 21376300 , PubMed: 26875626 , PubMed: 26919761 , PubMed: 27164704 , PubMed: 28095420 , PubMed: 28105280 , PubMed: 28126851 , PubMed: 28228639 ,

PubMed:[36959261](#), PubMed:[38538865](#), PubMed:[7679115](#), PubMed:[7681588](#), PubMed:[7685113](#)). NMDARs mediate simultaneously the potassium efflux and the influx of calcium and sodium (By similarity). Each GluN2 or GluN3 subunit confers differential attributes to channel properties, including activation, deactivation and desensitization kinetics, pH sensitivity, Ca²⁺ permeability, and binding to allosteric modulators (PubMed:[26875626](#), PubMed:[26919761](#), PubMed:[36309015](#), PubMed:[38598639](#)).


Cellular Location

Cell membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:P35439}. Postsynaptic cell membrane {ECO:0000250|UniProtKB:P35438}. Postsynaptic density membrane {ECO:0000250|UniProtKB:P35439}. Synaptic cell membrane {ECO:0000250|UniProtKB:P35438}. Note=Synaptic cell membrane targeting is dependent of GRIN2B/GluN2B subunit (By similarity). Association with GRIN3A occurs in the endoplasmic reticulum (By similarity) {ECO:0000250, ECO:0000250|UniProtKB:P35438, ECO:0000250|UniProtKB:P35439}

Images




Western blot analysis of NMDAR1 (Phospho-S890) expression in A549 (A), U87MG (B) whole cell lysates. (Predicted band size: 105 kD; Observed band size: 130 kD)



Immunohistochemical analysis of NMDAR1 (Phospho-S890) staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of NMDAR1 (Phospho-S890) staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a Alexa Fluor 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



Immunohistochemical analysis of NMDAR1 (pS890) staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.