

Anti-CDC25 (pS916) Antibody

Rabbit polyclonal antibody to CDC25 (pS916)

Catalog # AP61378

Product Information

Application	WB, IF/IC, IHC
Primary Accession	Q13972
Other Accession	P27671
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	145234

Additional Information

Gene ID	5923
Other Names	RASGRF1; CDC25; GNRP; GRF1; Ras-specific guanine nucleotide-releasing factor 1; Ras-GRF1; Guanine nucleotide-releasing protein; GNRP; Ras-specific nucleotide exchange factor CDC25
Target/Specificity	Recognizes endogenous levels of CDC25 (pS916) protein.
Dilution	WB~~WB (1/500 - 1/1000), IHC (1/50 - 1/200), IF/IC (1/100 - 1/500) IF/IC~~N/A IHC~~WB (1/500 - 1/1000), IHC (1/50 - 1/200), IF/IC (1/100 - 1/500)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt


Protein Information

Name	RASGRF1
Synonyms	CDC25, GNRP, GRF1
Function	Promotes the exchange of Ras-bound GDP by GTP.


Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human CDC25 (pS916). The exact sequence is proprietary.


Images



Western blot analysis of CDC25 (pS916) expression in U87MG (A), Panc1 (B), A549 (C) whole cell lysates.



Immunohistochemical analysis of CDC25 (pS916) staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of CDC25 (pS916) staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a Alexa Fluor 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.