

Anti-CD159c Antibody

Rabbit polyclonal antibody to CD159c

Catalog # AP61410

Product Information

Application	WB
Primary Accession	P26717
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	26159

Additional Information

Gene ID	3822
Other Names	NKG2C; NKG2-C type II integral membrane protein; CD159 antigen-like family member C; NK cell receptor C; NKG2-C-activating NK receptor; CD159c
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human CD159c. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	KLRC2
Synonyms	NKG2C {ECO:0000303 PubMed:18083576}
Function	Immune activating receptor involved in self-nonsel self discrimination. In complex with KLRD1 on cytotoxic lymphocyte subsets, recognizes non-classical major histocompatibility (MHC) class Ib HLA-E loaded with signal sequence-derived peptides from non-classical MHC class Ib HLA-G molecules, likely playing a role in the generation and effector functions of adaptive natural killer (NK) cells and in maternal-fetal tolerance during pregnancy (PubMed: 30134159 , PubMed: 37264229 , PubMed: 9754572). Regulates the effector functions of terminally differentiated cytotoxic lymphocyte subsets, and in particular may play a role in adaptive NK cell response to viral infection (PubMed: 20952657 , PubMed: 21825173). Upon HLA-E-peptide binding, transmits intracellular signals via the adapter protein TYROBP/DAP12, triggering the phosphorylation of proximal signaling molecules and cell

activation (PubMed:[15940674](#), PubMed:[9655483](#)).

Cellular Location

Cell membrane; Single-pass type II membrane protein


Tissue Location

Expressed in NK cell subsets, in particular in adaptive CD57-positive NK cells (at protein level) (PubMed:20952657, PubMed:21825173). Expressed in terminally differentiated cytotoxic gamma-delta T cells (at protein level) (PubMed:20952657). Expressed in alpha-beta T cells subsets (at protein level) (PubMed:20952657). KLRD1- KLRC1 and KLRD1-KLRC2 are differentially expressed within NK and T cell populations, with only minor subsets expressing both receptor complexes (at protein level) (PubMed:20952657).

Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human CD159c. The exact sequence is proprietary.

Images



Western blot analysis of CD159c expression in rat spleen (A), mouse spleen (B) whole cell lysates.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.