

# Anti-KANK2 Antibody

Rabbit polyclonal antibody to KANK2 Catalog # AP61469

#### **Product Information**

Application WB, IHC, IF
Primary Accession Q63ZY3
Reactivity Human
Host Rabbit
Clonality Polyclonal
Calculated MW 91174

#### **Additional Information**

**Gene ID** 25959

Other Names ANKRD25; KIAA1518; MXRA3; SIP; KN motif and ankyrin repeat

domain-containing protein 2; Ankyrin repeat domain-containing protein 25; Matrix-remodeling-associated protein 3; SRC-1-interacting protein; SIP;

SRC-interacting protein; SRC1-interacting protein

**Target/Specificity** KLH-conjugated synthetic peptide encompassing a sequence within the center

region of human KANK2. The exact sequence is proprietary.

**Dilution** WB~~WB (1/500 - 1/1000), IH (1/50 - 1/200), IF/IC (1/50 - 1/200)

IHC~~1:100~500 IF~~WB (1/500 - 1/1000), IH (1/50 - 1/200), IF/IC (1/50 -

1/200)

**Format** Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30%

glycerol, and 0.09% (W/V) sodium azide.

**Storage** Store at -20 °C.Stable for 12 months from date of receipt

#### **Protein Information**

Name KANK2

**Synonyms** ANKRD25, KIAA1518, MXRA3, SIP

**Function** Involved in transcription regulation by sequestering in the cytoplasm

nuclear receptor coactivators such as NCOA1, NCOA2 and NCOA3

(PubMed: <u>17476305</u>). Involved in regulation of caspase-independent apoptosis

by sequestering the proapoptotic factor AIFM1 in mitochondria (PubMed:22371500). Pro-apoptotic stimuli can induce its proteasomal degradation allowing the translocation of AIFM1 to the nucleus to induce apoptosis (PubMed:22371500). Involved in the negative control of vitamin D receptor signaling pathway (PubMed:24671081). Involved in actin stress fibers

formation through its interaction with ARHGDIA and the regulation of the Rho signaling pathway (PubMed:17996375, PubMed:25961457). May thereby play a role in cell adhesion and migration, regulating for instance podocytes migration during development of the kidney (PubMed:25961457). Through the Rho signaling pathway may also regulate cell proliferation (By similarity).

Cellular Location

Cytoplasm. Mitochondrion


**Tissue Location** 

Strongly expressed in cervix, colon, heart, kidney and lung. Expressed in kidney glomerular podocytes and mesangial cells (at protein level).


## **Background**

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human KANK2. The exact sequence is proprietary.


### **Images**



Western blot analysis of KANK2 expression in A375 (A), SKOVCAR3 (B) whole cell lysates.



Immunohistochemical analysis of KANK2 staining in human heart formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of KANK2 staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a Alexa Fluor 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.