

# Anti-Histone H2A (AcK9) Antibody

Rabbit polyclonal antibody to Histone H2A (AcK9) Catalog # AP61491

## **Product Information**

Bovine

## **Additional Information**

Gene ID	8329;8330;8332;8336;8969
Other Names	H2AFP; H2AFC; H2AFD; H2AFI; H2AFN; Histone H2A type 1; H2A.1; Histone H2A/p
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human Histone H2A with a site at AcK9. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt


#### **Protein Information**

Name	H2AC11 ( <u>HGNC:4737</u> )
Synonyms	H2AFP, HIST1H2AG
Function	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.
Cellular Location	Nucleus. Chromosome.

# Background

KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human Histone H2A with a site at AcK9. The exact sequence is proprietary.

#### Images



Western blot analysis of Histone H2A (AcK9) expression in Hela (A), H446 (B) whole cell lysates.

Direct ELISA antibody dose-response curve using Anti-Histone H2A (AcK9) Antibody. Antigen (acetyl-peptide and non-acetyl-peptide) concentration is 5 ug/ml. Goat Anti-Rabbit IgG (H&L) - HRP was used as the secondary antibody, and signal was developed by TMB substrate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.