

Anti-Histone H2A (AcK5) Antibody

Rabbit polyclonal antibody to Histone H2A (AcK5) Catalog # AP61492

Product Information

Application	WB
Primary Accession	<u>P0C0S8</u>
Other Accession	<u>P22752</u>
Reactivity	Human, Mouse, Rat, Chicken, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	14091
Reactivity Host Clonality	Human, Mouse, Rat, Chicken, Bovine Rabbit Polyclonal

Additional Information

Gene ID	8329;8330;8332;8336;8969
Other Names	H2AFP; H2AFC; H2AFD; H2AFI; H2AFN; Histone H2A type 1; H2A.1; Histone H2A/p
Target/Specificity	Recognizes endogenous levels of Histone H2A with a site at AcK5 protein.
Dilution	WB~~WB (1/500 - 1/1000)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	H2AC11 (<u>HGNC:4737</u>)
Synonyms	H2AFP, HIST1H2AG
Function	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.
Cellular Location	Nucleus. Chromosome.
Background	

KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human Histone H2A with a site at AcK5. The exact sequence is proprietary.

Images



Western blot analysis of Histone H2A (AcK5) expression in HEK293T (A), H446 (B), U2OS (C) whole cell lysates.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.