

Anti-EIF5A2 Antibody

Rabbit polyclonal antibody to EIF5A2

Catalog # AP61537

Product Information

Application	WB
Primary Accession	Q9GZV4
Other Accession	Q8BGY2
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	16793

Additional Information

Gene ID	56648
Other Names	Eukaryotic translation initiation factor 5A-2; eIF-5A-2; eIF-5A2; Eukaryotic initiation factor 5A isoform 2
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human EIF5A2. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	EIF5A2
Function	Translation factor that promotes translation elongation and termination, particularly upon ribosome stalling at specific amino acid sequence contexts (PubMed: 14622290). Binds between the exit (E) and peptidyl (P) site of the ribosome and promotes rescue of stalled ribosome: specifically required for efficient translation of polyproline-containing peptides as well as other motifs that stall the ribosome. Acts as a ribosome quality control (RQC) cofactor by joining the RQC complex to facilitate peptidyl transfer during CAT tailing step (By similarity). Also involved in actin dynamics and cell cycle progression, mRNA decay and probably in a pathway involved in stress response and maintenance of cell wall integrity (By similarity).
Cellular Location	Cytoplasm {ECO:0000250 UniProtKB:P63241}. Nucleus {ECO:0000250 UniProtKB:P63241}. Endoplasmic reticulum membrane

{ECO:0000250|UniProtKB:P63241}; Peripheral membrane protein
{ECO:0000250|UniProtKB:P63241}; Cytoplasmic side
{ECO:0000250|UniProtKB:P63241}. Note=Hypusine modification promotes the nuclear export and cytoplasmic localization and there was a dynamic shift in the localization from predominantly cytoplasmic to primarily nuclear under apoptotic inducing conditions {ECO:0000250|UniProtKB:P63241}


Tissue Location

Expressed in ovarian and colorectal cancer cell lines (at protein level). Highly expressed in testis. Overexpressed in some cancer cells.

Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human EIF5A2. The exact sequence is proprietary.

Images



Western blot analysis of EIF5A2 expression in AML12 (A), C6 (B), HEK293T (C), SGC7901 (D), HepG2 (E), A549 (F), A2780 (G) whole cell lysates.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.