

# Anti-IDE Antibody

Mouse monoclonal antibody to IDE

Catalog # AP61591

## Product Information

Application	WB
Primary Accession	<a href="#">P14735</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Calculated MW	117968

## Additional Information

Gene ID	3416
Other Names	Insulin-degrading enzyme; Abeta-degrading protease; Insulin protease; Insulinase; Insulysin
Target/Specificity	Recognizes endogenous levels of IDE protein.
Dilution	WB~~WB (1/1000 - 1/2000)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

## Protein Information

Name	IDE {ECO:0000303   PubMed:20364150, ECO:0000312   HGNC:HGNC:5381}
Function	Plays a role in the cellular breakdown of insulin, APP peptides, IAPP peptides, natriuretic peptides, glucagon, bradykinin, kallidin, and other peptides, and thereby plays a role in intercellular peptide signaling (PubMed: <a href="#">10684867</a> , PubMed: <a href="#">17051221</a> , PubMed: <a href="#">17613531</a> , PubMed: <a href="#">18986166</a> , PubMed: <a href="#">19321446</a> , PubMed: <a href="#">21098034</a> , PubMed: <a href="#">2293021</a> , PubMed: <a href="#">23922390</a> , PubMed: <a href="#">24847884</a> , PubMed: <a href="#">26394692</a> , PubMed: <a href="#">26968463</a> , PubMed: <a href="#">29596046</a> ). Substrate binding induces important conformation changes, making it possible to bind and degrade larger substrates, such as insulin (PubMed: <a href="#">23922390</a> , PubMed: <a href="#">26394692</a> , PubMed: <a href="#">29596046</a> ). Contributes to the regulation of peptide hormone signaling cascades and regulation of blood glucose homeostasis via its role in the degradation of insulin, glucagon and IAPP (By similarity). Plays a role in the degradation and clearance of APP-derived amyloidogenic peptides that are secreted by neurons and microglia (Probable) (PubMed: <a href="#">26394692</a> , PubMed: <a href="#">9830016</a> ). Degrades the natriuretic peptides ANP, BNP and CNP,

inactivating their ability to raise intracellular cGMP (PubMed:[21098034](#)). Also degrades an aberrant frameshifted 40-residue form of NPPA (fsNPPA) which is associated with familial atrial fibrillation in heterozygous patients (PubMed:[21098034](#)). Involved in antigen processing. Produces both the N terminus and the C terminus of MAGEA3-derived antigenic peptide (EVDPIGHLY) that is presented to cytotoxic T lymphocytes by MHC class I.

#### Cellular Location

Cytoplasm, cytosol. Cell membrane {ECO:0000250|UniProtKB:P35559}. Secreted Note=Present at the cell surface of neuron cells. The membrane-associated isoform is approximately 5 kDa larger than the known cytosolic isoform

#### Tissue Location

Detected in brain and in cerebrospinal fluid (at protein level).


## Background

---

KLH-conjugated synthetic peptide encompassing a sequence of human IDE. The exact sequence is proprietary.

## Images

---



Western blot analysis of IDE expression in Hela (A), HepG2 (B) whole cell lysates.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.