

Anti-IDE Antibody

Mouse monoclonal antibody to IDE

Catalog # AP61591

Product Information

Application	WB
Primary Accession	P14735
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Calculated MW	117968

Additional Information

Gene ID	3416
Other Names	Insulin-degrading enzyme; Abeta-degrading protease; Insulin protease; Insulinase; Insulysin
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence of human IDE. The exact sequence is proprietary.
Dilution	WB~~WB (1/1000 - 1/2000)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	IDE {ECO:0000303 PubMed:20364150, ECO:0000312 HGNC:HGNC:5381}
Function	Plays a role in the cellular breakdown of insulin, APP peptides, IAPP peptides, natriuretic peptides, glucagon, bradykinin, kallidin, and other peptides, and thereby plays a role in intercellular peptide signaling (PubMed: 10684867 , PubMed: 17051221 , PubMed: 17613531 , PubMed: 18986166 , PubMed: 19321446 , PubMed: 21098034 , PubMed: 2293021 , PubMed: 23922390 , PubMed: 24847884 , PubMed: 26394692 , PubMed: 26968463 , PubMed: 29596046). Substrate binding induces important conformation changes, making it possible to bind and degrade larger substrates, such as insulin (PubMed: 23922390 , PubMed: 26394692 , PubMed: 29596046). Contributes to the regulation of peptide hormone signaling cascades and regulation of blood glucose homeostasis via its role in the degradation of insulin, glucagon and IAPP (By similarity). Plays a role in the degradation and clearance of APP-derived amyloidogenic peptides that are secreted by neurons and microglia (Probable) (PubMed: 26394692 ,

PubMed:[9830016](#)). Degrades the natriuretic peptides ANP, BNP and CNP, inactivating their ability to raise intracellular cGMP (PubMed:[21098034](#)). Also degrades an aberrant frameshifted 40-residue form of NPPA (fsNPPA) which is associated with familial atrial fibrillation in heterozygous patients (PubMed:[21098034](#)). Involved in antigen processing. Produces both the N terminus and the C terminus of MAGEA3-derived antigenic peptide (EVDPIGHLY) that is presented to cytotoxic T lymphocytes by MHC class I.

Cellular Location

Cytoplasm, cytosol. Cell membrane {ECO:0000250|UniProtKB:P35559}. Secreted Note=Present at the cell surface of neuron cells. The membrane-associated isoform is approximately 5 kDa larger than the known cytosolic isoform


Tissue Location

Detected in brain and in cerebrospinal fluid (at protein level).

Background

KLH-conjugated synthetic peptide encompassing a sequence of human IDE. The exact sequence is proprietary.

Images



Western blot analysis of IDE expression in HeLa (A), HepG2 (B) whole cell lysates.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.