

Anti-MICU1 Antibody

Mouse monoclonal antibody to MICU1

Catalog # AP61592

Product Information

Application	WB, IHC
Primary Accession	Q9BPX6
Reactivity	Human, Mouse, Rat
Host	Mouse
Clonality	Monoclonal
Calculated MW	54351

Additional Information

Gene ID	10367
Other Names	CALC; CBARA1; Calcium uptake protein 1 mitochondrial; Atopy-related autoantigen CALC; ara CALC; Calcium-binding atopy-related autoantigen 1; allergen Hom s 4
Target/Specificity	Recombinant protein corresponding to human MICU1.
Dilution	WB~~WB (1/1000 - 1/2000), IHC (1/100 - 1/200) IHC~~WB (1/1000 - 1/2000), IHC (1/100 - 1/200)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	MICU1 {ECO:0000303 PubMed:20693986, ECO:0000312 HGNC:HGNC:1530}
Function	Calcium sensor of the mitochondrial calcium uniporter (MCU) channel, which senses calcium level via its EF-hand domains (PubMed: 20693986 , PubMed: 23101630 , PubMed: 23747253 , PubMed: 24313810 , PubMed: 24332854 , PubMed: 24503055 , PubMed: 24560927 , PubMed: 26341627 , PubMed: 26903221 , PubMed: 27099988 , PubMed: 28615291 , PubMed: 30454562 , PubMed: 30638448 , PubMed: 32494073 , PubMed: 32667285 , PubMed: 32762847 , PubMed: 32790952 , PubMed: 34463251 , PubMed: 36206740 , PubMed: 37036971 , PubMed: 37126688). MICU1 and MICU2 (or MICU3) form a disulfide-linked heterodimer that stimulates and inhibits MCU activity, depending on the concentration of calcium (PubMed: 24560927 , PubMed: 26903221 , PubMed: 28615291 , PubMed: 32148862 , PubMed: 32494073 , PubMed: 32667285 , PubMed: 32762847 ,

PubMed:[32790952](#), PubMed:[36206740](#), PubMed:[37036971](#), PubMed:[37126688](#)). At low calcium levels, MICU1 occludes the pore of the MCU channel, preventing mitochondrial calcium uptake (PubMed:[32494073](#), PubMed:[32667285](#), PubMed:[32762847](#), PubMed:[37036971](#), PubMed:[37126688](#)). At higher calcium levels, calcium-binding to MICU1 and MICU2 (or MICU3) induces a conformational change that weakens MCU-MICU1 interactions and moves the MICU1-MICU2 heterodimer away from the pore, allowing calcium permeation through the MCU channel (PubMed:[32494073](#), PubMed:[32667285](#), PubMed:[32762847](#)). Also required to protect against manganese toxicity by preventing manganese uptake by MCU: mechanistically, manganese-binding to its EF-hand domains does not induce any conformational change, maintaining MCU pore occlusion (PubMed:[30082385](#), PubMed:[30403999](#)). Also acts as a barrier for inhibitors of the MCU channel, such as ruthenium red or its derivative Ru360 (PubMed:[37244260](#)). Acts as a regulator of mitochondrial cristae structure independently of its ability to regulate the mitochondrial calcium uniporter channel (PubMed:[31427612](#), PubMed:[37098122](#)). Regulates glucose-dependent insulin secretion in pancreatic beta-cells by regulating mitochondrial calcium uptake (PubMed:[22904319](#)). Induces T-helper 1-mediated autoreactivity, which is accompanied by the release of IFNG (PubMed:[16002733](#)).

Cellular Location

Mitochondrion intermembrane space. Mitochondrion inner membrane. Note=Recruited to the mitochondrial inner membrane by EMRE/SMDT1 (PubMed:30454562). Also localizes to mitochondrial cristae junctions (PubMed:31427612)


Tissue Location

Expressed in epithelial cell lines. Strongly expressed in epidermal keratinocytes and dermal endothelial cells

Background


Recombinant protein corresponding to human MICU1.

Images



Western blot analysis of MICU1 expression in MCF7 (A), mouse brain (B), rat brain (C) whole cell lysates.

Immunohistochemical analysis of MICU1 staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.