

# Anti-COVID-19 Spike glycoprotein Antibody

Rabbit polyclonal antibody to COVID-19 Spike glycoprotein

Catalog # AP61626

## Product Information

Application	WB, E
Primary Accession	<a href="#">P0DTC2</a>
Host	Rabbit
Clonality	Polyclonal
Calculated MW	141178

## Additional Information

Gene ID	43740568
Other Names	Spike glycoprotein; S glycoprotein; E2; Peplomer protein
Target/Specificity	Recognizes COVID-19 Spike glycoprotein.
Dilution	WB~~1:1000 E~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

## Protein Information

Name	S {ECO:0000255 HAMAP-Rule:MF_04099}
Function	[Spike protein S1]: Attaches the virion to the cell membrane by interacting with host receptor, initiating the infection. The major receptor is host ACE2 (PubMed: <a href="#">32142651</a> , PubMed: <a href="#">32155444</a> , PubMed: <a href="#">33607086</a> ). When S2/S2' has been cleaved, binding to the receptor triggers direct fusion at the cell membrane (PubMed: <a href="#">34561887</a> ). When S2/S2' has not been cleaved, binding to the receptor results in internalization of the virus by endocytosis using host TFRC and GRM2 and leading to fusion of the virion membrane with the host endosomal membrane (PubMed: <a href="#">32075877</a> , PubMed: <a href="#">32221306</a> , PubMed: <a href="#">34903715</a> , PubMed: <a href="#">36779763</a> ). Alternatively, may use NRP1/NRP2 (PubMed: <a href="#">33082294</a> , PubMed: <a href="#">33082293</a> ) and integrin as entry receptors (PubMed: <a href="#">35150743</a> ). The use of NRP1/NRP2 receptors may explain the tropism of the virus in human olfactory epithelial cells, which express these molecules at high levels but ACE2 at low levels (PubMed: <a href="#">33082293</a> ). The stalk domain of S contains three hinges, giving the head unexpected orientational freedom (PubMed: <a href="#">32817270</a> ).
Cellular Location	Virion membrane {ECO:0000255 HAMAP-Rule:MF_04099,

ECO:0000269 | PubMed:32979942}; Single-pass type I membrane protein {ECO:0000255 | HAMAP-Rule:MF\_04099, ECO:0000269 | PubMed:34504087}. Host endoplasmic reticulum-Golgi intermediate compartment membrane {ECO:0000255 | HAMAP-Rule:MF\_04099, ECO:0000269 | PubMed:34504087}; Single-pass type I membrane protein {ECO:0000255 | HAMAP-Rule:MF\_04099}. Host cell membrane {ECO:0000255 | HAMAP-Rule:MF\_04099, ECO:0000269 | PubMed:34504087}; Single-pass type I membrane protein {ECO:0000255 | HAMAP-Rule:MF\_04099}. Note=Accumulates in the endoplasmic reticulum-Golgi intermediate compartment, where it participates in virus particle assembly. Some S oligomers are transported to the host plasma membrane, where they may mediate cell-cell fusion (PubMed:34504087). An average of 26 +/-15 S trimers are found randomly distributed at the surface of the virion (PubMed:32979942) {ECO:0000255 | HAMAP-Rule:MF\_04099, ECO:0000269 | PubMed:32979942, ECO:0000269 | PubMed:34504087}

## Background

---

KLH-conjugated synthetic peptide encompassing a sequence within the center region of COVID-19 Spike glycoprotein. The exact sequence is proprietary.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.