

Cleaved-Cathepsin C HC (R394) Polyclonal Antibody

Catalog # AP63110

Product Information

Application	WB, E
Primary Accession	P53634
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	51854

Additional Information

Gene ID	1075
Other Names	CTSC; CPPI; Dipeptidyl peptidase 1; Cathepsin C; Cathepsin J; Dipeptidyl peptidase I; DPP-I; DPPI; Dipeptidyl transferase
Dilution	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications. E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	CTSC
Synonyms	CPPI
Function	Thiol protease (PubMed: 1586157). Has dipeptidylpeptidase activity (PubMed: 1586157). Active against a broad range of dipeptide substrates composed of both polar and hydrophobic amino acids (PubMed: 1586157). Proline cannot occupy the P1 position and arginine cannot occupy the P2 position of the substrate (PubMed: 1586157). Can act as both an exopeptidase and endopeptidase (PubMed: 1586157). Activates serine proteases such as elastase, cathepsin G and granzymes A and B (PubMed: 8428921).
Cellular Location	Lysosome.
Tissue Location	Ubiquitous. Highly expressed in lung, kidney and placenta. Detected at intermediate levels in colon, small intestine, spleen and pancreas.

Background

Thiol protease. Has dipeptidylpeptidase activity. Active against a broad range of dipeptide substrates composed of both polar and hydrophobic amino acids. Proline cannot occupy the P1 position and arginine cannot occupy the P2 position of the substrate. Can act as both an exopeptidase and endopeptidase. Activates serine proteases such as elastase, cathepsin G and granzymes A and B. Can also activate neuraminidase and factor XIII.

Images



Western Blot analysis of various cells using
Cleaved-Cathepsin C HC (R394) Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.