

# Cleaved-Notch 2 (A1734) Polyclonal Antibody

Catalog # AP63143

#### **Product Information**

Application WB Primary Accession Q04721

Reactivity Human, Mouse, Rat

HostRabbitClonalityPolyclonalCalculated MW265405

#### **Additional Information**

**Gene ID** 4853

Other Names NOTCH2; Neurogenic locus notch homolog protein 2; Notch 2; hN2

Dilution WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other

applications.

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

## **Protein Information**

Name NOTCH2 ( HGNC:7882)

**Function** Functions as a receptor for membrane-bound ligands Jagged-1 (JAG1),

Jagged-2 (JAG2) and Delta-1 (DLL1) to regulate cell-fate determination. Upon ligand activation through the released notch intracellular domain (NICD) it forms a transcriptional activator complex with RBPJ/RBPSUH and activates genes of the enhancer of split locus (PubMed:21378985, PubMed:21378989). Affects the implementation of differentiation, proliferation and apoptotic programs (By similarity). Involved in bone remodeling and homeostasis. In collaboration with RELA/p65 enhances NFATc1 promoter activity and

positively regulates RANKL-induced osteoclast differentiation

(PubMed: <u>29149593</u>). Positively regulates self-renewal of liver cancer cells

(PubMed: 25985737).

**Cellular Location** [Notch 2 extracellular truncation]: Cell membrane; Single-pass type I

membrane protein


**Tissue Location** Expressed in the brain, heart, kidney, lung, skeletal muscle and liver.

Ubiquitously expressed in the embryo

## **Background**

Functions as a receptor for membrane-bound ligands Jagged-1 (JAG1), Jagged-2 (JAG2) and Delta-1 (DLL1) to regulate cell-fate determination. Upon ligand activation through the released notch intracellular domain (NICD) it forms a transcriptional activator complex with RBPJ/RBPSUH and activates genes of the enhancer of split locus (PubMed:21378985, PubMed:21378989). Affects the implementation of differentiation, proliferation and apoptotic programs (By similarity). Involved in bone remodeling and homeostasis. In collaboration with RELA/p65 enhances NFATc1 promoter activity and positively regulates RANKL- induced osteoclast differentiation (PubMed:29149593). Positively regulates self-renewal of liver cancer cells (PubMed:25985737).

## **Images**



Western Blot analysis of Jurkat cells using Cleaved-Notch 2 (A1734) Polyclonal Antibody diluted at 1: 2000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.