

Cleaved-Thrombin APII (R327) Polyclonal Antibody

Catalog # AP63152

Product Information

Application	WB
Primary Accession	P00734
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	70037

Additional Information

Gene ID	2147
Other Names	F2; Prothrombin; Coagulation factor II
Dilution	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	F2
Function	Thrombin, which cleaves bonds after Arg and Lys, converts fibrinogen to fibrin and activates factors V, VII, VIII, XIII, and, in complex with thrombomodulin, protein C. Functions in blood homeostasis, inflammation and wound healing. Activates coagulation factor XI (F11); activation is promoted by the contact with negatively charged surfaces (PubMed: 2019570 , PubMed: 21976677). Triggers the production of pro- inflammatory cytokines, such as MCP-1/CCL2 and IL8/CXCL8, in endothelial cells (PubMed: 30568593 , PubMed: 9780208).
Cellular Location	Secreted, extracellular space.
Tissue Location	Expressed by the liver and secreted in plasma.

Background

Thrombin, which cleaves bonds after Arg and Lys, converts fibrinogen to fibrin and activates factors V, VII, VIII, XIII, and, in complex with thrombomodulin, protein C. Functions in blood homeostasis, inflammation

and wound healing.

Images



Western Blot analysis of various cells using
Cleaved-Thrombin APII (R327) Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.