

# Cleaved-Factor X/ Factor X LC (A41) Polyclonal Antibody

Catalog # AP63157

## Product Information

---

Application	WB
Primary Accession	<a href="#">P00742</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	54732

## Additional Information

---

Gene ID	2159
Other Names	F10; Coagulation factor X; Stuart factor; Stuart-Prower factor
Dilution	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

## Protein Information

---

Name	F10
Function	<p>Factor Xa is a vitamin K-dependent glycoprotein that converts prothrombin to thrombin in the presence of factor Va, calcium and phospholipid during blood clotting (PubMed:<a href="#">22409427</a>). Factor Xa activates pro-inflammatory signaling pathways in a protease-activated receptor (PAR)-dependent manner (PubMed:<a href="#">24041930</a>, PubMed:<a href="#">30568593</a>, PubMed:<a href="#">34831181</a>, PubMed:<a href="#">18202198</a>). Up-regulates expression of protease- activated receptors (PARs) F2R, F2RL1 and F2RL2 in dermal microvascular endothelial cells (PubMed:<a href="#">35738824</a>). Triggers the production of pro- inflammatory cytokines, such as MCP-1/CCL2 and IL6, in cardiac fibroblasts and umbilical vein endothelial cells in PAR-1/F2R-dependent manner (PubMed:<a href="#">30568593</a>, PubMed:<a href="#">34831181</a>). Triggers the production of pro-inflammatory cytokines, such as MCP-1/CCL2, IL6, TNF-alpha/TNF, IL- 1beta/IL1B, IL8/CXCL8 and IL18, in endothelial cells and atrial tissues (PubMed:<a href="#">24041930</a>, PubMed:<a href="#">35738824</a>, PubMed:<a href="#">9780208</a>). Induces expression of adhesion molecules, such as ICAM1, VCAM1 and SELE, in endothelial cells and atrial tissues (PubMed:<a href="#">24041930</a>, PubMed:<a href="#">35738824</a>, PubMed:<a href="#">9780208</a>). Increases expression of phosphorylated ERK1/2 in dermal microvascular endothelial cells and atrial tissues (PubMed:<a href="#">24041930</a>, PubMed:<a href="#">35738824</a>). Triggers activation of the</p>

transcription factor NF-kappa-B in dermal microvascular endothelial cells and atrial tissues (PubMed:[24041930](#), PubMed:[35738824](#)). Activates pro-inflammatory and pro-fibrotic responses in dermal fibroblasts and enhances wound healing probably via PAR-2/F2RL1-dependent mechanism (PubMed:[18202198](#)). Activates barrier protective signaling responses in endothelial cells in PAR-2/F2RL1-dependent manner; the activity depends on the cleavage of PAR-2/F2RL1 by factor Xa (PubMed:[22409427](#)). Up-regulates expression of plasminogen activator inhibitor 1 (SERPINE1) in atrial tissues (PubMed:[24041930](#)).

**Cellular Location**

Secreted.

**Tissue Location**

Plasma; synthesized in the liver.


## Background

---

Factor Xa is a vitamin K-dependent glycoprotein that converts prothrombin to thrombin in the presence of factor Va, calcium and phospholipid during blood clotting.

## Images

---



Western Blot analysis of various cells using  
Cleaved-Factor X/ Factor X LC (A41) Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.