

Cleaved-Factor XII HC (R372) Polyclonal Antibody

Catalog # AP63160

Product Information

Application	WB
Primary Accession	P00748
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	67792

Additional Information

Gene ID	2161
Other Names	F12; Coagulation factor XII; Hageman factor; HAF
Dilution	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C


Protein Information

Name	F12
Function	Factor XII is a serum glycoprotein that participates in the initiation of blood coagulation, fibrinolysis, and the generation of bradykinin and angiotensin. Prekallikrein is cleaved by factor XII to form kallikrein, which then cleaves factor XII first to alpha-factor XIIa and then trypsin cleaves it to beta-factor XIIa. Alpha-factor XIIa activates factor XI to factor XIa (PubMed: 2019570 , PubMed: 21304106 , PubMed: 8427954).
Cellular Location	Secreted.

Background

Factor XII is a serum glycoprotein that participates in the initiation of blood coagulation, fibrinolysis, and the generation of bradykinin and angiotensin. Prekallikrein is cleaved by factor XII to form kallikrein, which then cleaves factor XII first to alpha-factor XIIa and then trypsin cleaves it to beta- factor XIIa. Alpha-factor XIIa activates factor XI to factor XIa.

Images



Western Blot analysis of various cells using
Cleaved-Factor XII HC (R372) Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.