

# Cleaved-MPO 89k (A49) Polyclonal Antibody

Catalog # AP63175

## Product Information

---

<b>Application</b>	WB
<b>Primary Accession</b>	<a href="#">P05164</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	83869

## Additional Information

---

<b>Gene ID</b>	4353
<b>Other Names</b>	MPO; Myeloperoxidase; MPO
<b>Dilution</b>	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
<b>Format</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
<b>Storage Conditions</b>	-20°C

## Protein Information


---

<b>Name</b>	MPO ( <a href="#">HGNC:7218</a> )
<b>Function</b>	Part of the host defense system of polymorphonuclear leukocytes. It is responsible for microbicidal activity against a wide range of organisms. In the stimulated PMN, MPO catalyzes the production of hypohalous acids, primarily hypochlorous acid in physiologic situations, and other toxic intermediates that greatly enhance PMN microbicidal activity (PubMed: <a href="#">9922160</a> ). Mediates the proteolytic cleavage of alpha-1-microglobulin to form t-alpha-1-microglobulin, which potently inhibits oxidation of low-density lipoprotein particles and limits vascular damage (PubMed: <a href="#">25698971</a> ).
<b>Cellular Location</b>	Lysosome.

## Background

---

Part of the host defense system of polymorphonuclear leukocytes. It is responsible for microbicidal activity against a wide range of organisms. In the stimulated PMN, MPO catalyzes the production of hypohalous acids, primarily hypochlorous acid in physiologic situations, and other toxic intermediates that greatly enhance PMN microbicidal activity.



Western Blot analysis of various cells using Cleaved-MPO 89k (A49) Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.