

# Histone H3 (Mono Methyl Lys80) Polyclonal Antibody

Catalog # AP63188

### **Product Information**

**Application** WB

Primary Accession P68431, Q71DI3, P84243, Q6NXT2

Reactivity Human, Mouse, Rat

Host Rabbit
Clonality Polyclonal
Calculated MW 15404

## **Additional Information**

**Gene ID** 8350;8351;8352;8353;8354;8355;8356;8357;8358;8968

Other Names HIST1H3A; H3FA; HIST1H3B; H3FL; HIST1H3C; H3FC; HIST1H3D; H3FB;

HIST1H3E; H3FD; HIST1H3F; H3FI; HIST1H3G; H3FH; HIST1H3H; H3FK; HIST1H3I; H3FF; HIST1H3J; H3FJ; Histone H3.1; Histone H3.1; Histone H3.2; Histone H3/m; Histone H3/o; H3F3A; H3.3A; H3F3; PP781; H3F3B; H3.3B;

Histone H3.3; H3F3C; Histone H3.3C; Histone H3.5

**Dilution** WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other

applications.

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

### **Protein Information**

Name H3C1 ( <u>HGNC:4766</u>)

**Synonyms** H3FA, HIST1H3A


**Function** Core component of nucleosome. Nucleosomes wrap and compact DNA into

chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications

of histones, also called histone code, and nucleosome remodeling.

**Cellular Location** Nucleus. Chromosome.

# **Images**



Western Blot analysis of MCF-7 cells using Mono-Methyl-Histone H3 (K80) Polyclonal Antibody.. Secondary antibody was diluted at 1:20000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.