

# Tubulin α (Acetyl Lys112) Polyclonal Antibody

Catalog # AP63242

#### **Product Information**

**Application** WB

Primary Accession Q71U36, P68363, Q9BQE3
Reactivity Human, Mouse, Rat

Host Rabbit
Clonality Polyclonal
Calculated MW 50136

### **Additional Information**

**Gene ID** 7846

Other Names TUBA1A; TUBA3; Tubulin alpha-1A chain; Alpha-tubulin 3; Tubulin B-alpha-1;

Tubulin alpha-3 chain; TUBA1B; Tubulin alpha-1B chain; Alpha-tubulin ubiquitousTubulin K-alpha-1; Tubulin alpha-ubiquitous chain; TUBA1C; TUBA6; Tubulin alpha-1C chain; Alpha-tubulin 6; Tubulin alpha-6 chain

Dilution WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other

applications.

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

#### **Protein Information**

Name TUBA1A

Synonyms TUBA3

**Function** Tubulin is the major constituent of microtubules, a cylinder consisting of

laterally associated linear protofilaments composed of alpha- and

beta-tubulin heterodimers. Microtubules grow by the addition of GTP-tubulin dimers to the microtubule end, where a stabilizing cap forms. Below the cap,

tubulin dimers are in GDP-bound state, owing to GTPase activity of


alpha-tubulin.

**Cellular Location** Cytoplasm, cytoskeleton, flagellum axoneme

{ECO:0000250 | UniProtKB:P68369}

**Tissue Location** Expressed at a high level in fetal brain.

## **Images**



Western Blot analysis of HepG2 cells using Acetyl-Tubulin  $\alpha$  (K112) Polyclonal Antibody.. Secondary antibody was diluted at 1:20000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.