

# Aquaporin 4 Monoclonal Antibody(4H1)

Catalog # AP63316

## Product Information

---

Application	WB, IHC-P, IF
Primary Accession	<a href="#">P55087</a>
Reactivity	Human, Mouse, Rat
Host	Mouse
Clonality	Monoclonal
Calculated MW	34830

## Additional Information

---

Gene ID	361
Other Names	AQP4; Aquaporin-4; AQP-4; Mercurial-insensitive water channel; MIWC; WCH4
Dilution	WB~~WB: 1:1000 IF: 1:100-200 IHC 1:50-300 IHC-P~~N/A IF~~1:50~200
Format	PBS, pH 7.4, containing 0.09% (W/V) sodium azide as Preservative and 50% Glycerol.
Storage Conditions	-20°C

## Protein Information

---

Name	AQP4
Function	Forms a water-specific channel (PubMed: <a href="#">19383790</a> , PubMed: <a href="#">7559426</a> , PubMed: <a href="#">8601457</a> ). Plays an important role in brain water homeostasis (PubMed: <a href="#">37143309</a> ). It is involved in glymphatic solute transport and is required for a normal rate of water exchange across the blood brain interface. Required for normal levels of cerebrospinal fluid influx into the brain cortex and parenchyma along paravascular spaces that surround penetrating arteries, and for normal drainage of interstitial fluid along paravenous drainage pathways. Thereby, it is required for normal clearance of solutes from the brain interstitial fluid, including soluble beta-amyloid peptides derived from APP. Plays a redundant role in urinary water homeostasis and urinary concentrating ability (By similarity).
Cellular Location	Cell membrane; Multi-pass membrane protein. Basolateral cell membrane {ECO:0000250 UniProtKB:P55088}; Multi-pass membrane protein. Endosome membrane {ECO:0000250 UniProtKB:P47863}. Cell membrane, sarcolemma; Multi-pass membrane protein. Cell projection {ECO:0000250 UniProtKB:P47863}. Note=Activation of the vasopressin receptor AVPR1A triggers AQP4 phosphorylation at Ser-180 and promotes its internalization from the cell membrane. Detected on brain astrocyte

processes and astrocyte endfeet close to capillaries  
{ECO:0000250|UniProtKB:P47863}


## Tissue Location

Detected in skeletal muscle (PubMed:29055082). Detected in stomach, along the glandular base region of the fundic gland (at protein level) (PubMed:8601457). Detected in brain, lung and skeletal muscle, and at much lower levels in heart and ovary (PubMed:7559426, PubMed:8601457).


## Background

Forms a water-specific channel. Osmoreceptor which regulates body water balance and mediates water flow within the central nervous system.


## Images




Immunohistochemical analysis of paraffin-embedded Human-liver-cancer tissue. 1, Aquaporin 4 Monoclonal Antibody(4H1) was diluted at 1:200(4°C, overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval(>98°C, 20min). 3, Secondary antibody was diluted at 1:200(room temperature, 30min). Negative control was used by secondary antibody only.




Immunofluorescence analysis of Human-appendix tissue. 1, Aquaporin 4 Monoclonal Antibody(4H1)(red) was diluted at 1:200(4°C, overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300(room temperature, 50min). 3, Picture B: DAPI(blue) 10min. Picture A: Target. Picture B: DAPI. Picture C: merge of A+B



Immunofluorescence analysis of Mouse-brain tissue. 1, Aquaporin 4 Monoclonal Antibody(4H1)(red) was diluted at 1:200(4°C, overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300(room temperature, 50min). 3, Picture B: DAPI(blue) 10min. Picture A: Target. Picture B: DAPI. Picture C: merge of A+B



Western blot analysis of 1) HeLa, 2) Mouse Heart tissue, 3) Rat Heart Tissue, diluted at 1:2000.



IF analysis of HeLa with antibody (Left) and DAPI (Right) diluted at 1:100.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.