

# CD10 Monoclonal Antibody(5B8)

Catalog # AP63326

## Product Information

Application	IHC-P, IF
Primary Accession	<a href="#">P08473</a>
Reactivity	Human, Mouse, Rat
Host	Mouse
Clonality	Monoclonal
Calculated MW	85514

## Additional Information

Gene ID	4311
Other Names	MME; EPN; Neprilysin; Atriopeptidase; Common acute lymphocytic leukemia antigen; CALLA; Enkephalinase; Neutral endopeptidase 24.11; NEP; Neutral endopeptidase; Skin fibroblast elastase; SFE; CD10
Dilution	IHC-P~~N/A IF~~IF: 1:50-200 IHC: 1:200
Format	PBS, pH 7.4, containing 0.09% (W/V) sodium azide as Preservative and 50% Glycerol.
Storage Conditions	-20°C


## Protein Information

Name	MME {ECO:0000303 PubMed:27588448, ECO:0000312 HGNC:HGNC:7154}
Function	Thermolysin-like specificity, but is almost confined on acting on polypeptides of up to 30 amino acids (PubMed: <a href="#">15283675</a> , PubMed: <a href="#">6208535</a> , PubMed: <a href="#">6349683</a> , PubMed: <a href="#">8168535</a> ). Biologically important in the destruction of opioid peptides such as Met- and Leu-enkephalins by cleavage of a Gly-Phe bond (PubMed: <a href="#">17101991</a> , PubMed: <a href="#">6349683</a> ). Catalyzes cleavage of bradykinin, substance P and neurotensin peptides (PubMed: <a href="#">6208535</a> ). Able to cleave angiotensin-1, angiotensin-2 and angiotensin 1-9 (PubMed: <a href="#">15283675</a> , PubMed: <a href="#">6349683</a> ). Involved in the degradation of atrial natriuretic factor (ANF) and brain natriuretic factor (BNP(1-32)) (PubMed: <a href="#">16254193</a> , PubMed: <a href="#">2531377</a> , PubMed: <a href="#">2972276</a> ). Displays UV-inducible elastase activity toward skin preelastic and elastic fibers (PubMed: <a href="#">20876573</a> ).
Cellular Location	Cell membrane; Single-pass type II membrane protein


## Background

Thermolysin-like specificity, but is almost confined on acting on polypeptides of up to 30 amino acids (PubMed:[15283675](#), PubMed:[8168535](#)). Biologically important in the destruction of opioid peptides such as Met- and Leu-enkephalins by cleavage of a Gly-Phe bond (PubMed:[17101991](#)). Able to cleave angiotensin-1, angiotensin-2 and angiotensin 1-9 (PubMed:[15283675](#)). Involved in the degradation of atrial natriuretic factor (ANF) (PubMed:[2531377](#), PubMed:[2972276](#)). Displays UV-inducible elastase activity toward skin preelastic and elastic fibers (PubMed:[20876573](#)).


## Images




Immunofluorescence analysis of human-uterus tissue. 1,CD10 Monoclonal Antibody(5B8)(red) was diluted at 1:200(4°C,overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300(room temperature, 50min).3, Picture B: DAPI(blue) 10min. Picture A:Target. Picture B: DAPI. Picture C: merge of A+B




Immunofluorescence analysis of rat-kidney tissue. 1,CD10 Monoclonal Antibody(5B8)(red) was diluted at 1:200(4°C,overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300(room temperature, 50min).3, Picture B: DAPI(blue) 10min. Picture A:Target. Picture B: DAPI. Picture C: merge of A+B



Immunohistochemical analysis of paraffin-embedded Human-uterus tissue. 1,CD10 Monoclonal Antibody(5B8) was diluted at 1:200(4°C,overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200(room temperature, 30min). Negative control was used by secondary antibody only.




Immunohistochemical analysis of paraffin-embedded Rat-kidney tissue. 1,CD10 Monoclonal Antibody(5B8) was diluted at 1:200(4°C,overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200(room temperature, 30min). Negative control was used by secondary antibody only.



Immunohistochemical analysis of paraffin-embedded Mouse-lung tissue. 1,CD10 Monoclonal Antibody(5B8) was diluted at 1:200(4°C,overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200(room temperature, 30min). Negative control was used by secondary antibody only.

Immunohistochemical analysis of paraffin-embedded human-breast-cancer using antibody diluted at 1:50.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.