

Histone H3 (Tri Methyl Lys9) Monoclonal Antibody(2C3)

Catalog # AP63354

Product Information

Application WB

Primary Accession P68431, Q71DI3, P84243
Reactivity Human, Mouse, Rat

Host Mouse
Clonality Monoclonal
Calculated MW 15404

Additional Information

Gene ID 8350;8351;8352;8353;8354;8355;8356;8357;8358;8968

Other Names HIST1H3A; H3FA; HIST1H3B; H3FL; HIST1H3C; H3FC; HIST1H3D; H3FB;

HIST1H3E; H3FD; HIST1H3F; H3FI; HIST1H3G; H3FH; HIST1H3H; H3FK; HIST1H3I; H3FF; HIST1H3J; H3FJ; Histone H3.1; Histone H3/a; Histone H3/b; Histone H3/c; Histone H3/d; Histone H3/f; Histone H3/h; Histone H3/i; Histone

H3/j; Histone H3/k; Histone H3/l; HIST2H3A; HIST2H3C; H3F2; H3FM;

HIST2H3D; Histone H3.2; Histone H3/m; Histone H3/o; H3F3A; H3.3A; H3F3;

PP781; H3F3B; H3.3B; Histone H3.3

Dilution WB~~WB: 1:1000-2000

Format PBS, pH 7.4, containing 0.09% (W/V) sodium azide as Preservative and 50%

Glycerol.

Storage Conditions -20°C

Protein Information

Name H3C1 (<u>HGNC:4766</u>)

Synonyms H3FA, HIST1H3A


Function Core component of nucleosome. Nucleosomes wrap and compact DNA into

chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications


of histones, also called histone code, and nucleosome remodeling.

Cellular Location Nucleus. Chromosome.

Images



Western blot analysis of 1) Hela, 2) Rat Testis tissue, 3) Raw264.7, diluted at 1:2000. cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).



The picture was kindly provided by our customer

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.