

# Histone H4 (Mono Methyl Lys79) Polyclonal Antibody

Catalog # AP63427

## **Product Information**

Application WB Primary Accession P62805

**Reactivity** Human, Mouse, Rat

HostMouseClonalityMonoclonalCalculated MW11367

### **Additional Information**

**Gene ID** 121504;554313;8294;8359;8360;8361;8362;8363;8364;8365;8366;8367;8368;

8370

Other Names HIST1H4A; H4/A; H4FA; HIST1H4B; H4/I; H4FI; HIST1H4C; H4/G; H4FG;

HIST1H4D; H4/B; H4FB; HIST1H4E; H4/J; H4FJ; HIST1H4F; H4/C; H4FC; HIST1H4H; H4/H; H4FH; HIST1H4I; H4/M; H4FM; HIST1H4J; H4/E; H4FE; HIST1H4K; H4/D; H4FD; HIST1H4L; H4/K; H4FK; HIST2H4A; H4/N; H4F2; H4FN;

HIST2H4; HIST2H4B; H4/O; H4FO; HIST4H4; Histone H4

**Dilution** WB~~WB: 1:500-1000

**Format** PBS, pH 7.4, containing 0.09% (W/V) sodium azide as Preservative and 50%

Glycerol.

Storage Conditions -20°C

#### **Protein Information**

Name H4C1

Synonyms H4/A, H4FA, HIST1H4A

**Function** Core component of nucleosome. Nucleosomes wrap and compact DNA into

chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications

of histones, also called histone code, and nucleosome remodeling.

Cellular Location Nucleus {ECO:0000250 | UniProtKB:P62806}. Chromosome. Note=Localized to


the nucleus when acetylated in step 11 spermatids.

{ECO:0000250 | UniProtKB:P62806}

# **Background**

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

# **Images**



Western blot analysis of 1) Hela, 2) 3T3, diluted at 1:2000.. Secondary antibody was diluted at 1:20000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.